

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 September 2003 (12.09.2003)


PCT

(10) International Publication Number
WO 03/074520 A1

- (51) International Patent Classification⁷: **C07D 417/04**,
A61K 31/496, A61P 25/22, 25/24, 29/00 // (C07D 417/04,
241:00, 285:00)
- (74) Agents: **ABRAMS, Samuel, B.** et al.; Pennie & Edmonds
L.L.P., 1155 Avenue of the Americas, New York, NY 10036
(US).
- (21) International Application Number: **PCT/US03/06107**
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 28 February 2003 (28.02.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/360,172 1 March 2002 (01.03.2002) US
60/411,084 17 September 2002 (17.09.2002) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **EURO-
CELTIQUE, S.A.** [LU/LU]; 122 Boulevard de la Petrusse,
L-2230 Luxembourg (LU).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **KYLE, Donald,
J.** [US/US]; 29 Weatherfield Drive, Newtown, PA 18940
(US). **SUN, Qun** [CN/US]; 108 Wrangle Court, #6,
Princeton, NJ 08540 (US).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

BEST AVAILABLE COPY

(54) Title: **THIADIAZOLYLPIPERAZINE DERIVATIVES USEFUL FOR TREATING OR PREVENTING PAIN**



(57) Abstract: A compound of formula (I) (wherein X is O or S and R¹, R³, R⁴, and R⁵ are disclosed herein) or a pharmaceutically acceptable salt thereof (a "Thiadiazolylpiperazine Compound"), pharmaceutical compositions comprising a Thiadiazolylpiperazine Compound, and methods for treating or preventing pain in a patient comprising administering to a patient in need thereof an effective amount of a Thiadiazolylpiperazine Compound are disclosed.

WO 03/074520 A1

THIADIAZOLYLPIPERAZINE DERIVATIVES USEFUL FOR TREATING OR PREVENTING PAIN

This application claims the benefit of U.S. Provisional Application No. 60/360,172, filed March 1, 2002, and U.S. Provisional Application No. 60/411,084, filed September 17, 2002, which are incorporated herein by reference in their entirety.

1. FIELD OF THE INVENTION

The present invention relates to Thiadiazolylpiperazine Compounds; compositions comprising a Thiadiazolylpiperazine Compound; and methods for preventing or treating pain, urinary incontinence (UI), an ulcer, inflammatory-bowel disease (IBD), irritable-bowel syndrome (IBS), an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, amyotrophic lateral sclerosis (ALS), dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia or depression, comprising administering to an animal in need thereof an effective amount of a Thiadiazolylpiperazine Compound.

2. BACKGROUND OF THE INVENTION

Pain is the most common symptom for which patients seek medical advice and treatment. Pain can be acute or chronic. While acute pain is usually self-limited, chronic pain persists for 3 months or longer and can lead to significant changes in a patient's personality, lifestyle, functional ability and overall quality of life (K.M. Foley, Pain, in *Cecil Textbook of Medicine* 100-107 (J.C. Bennett and F. Plum eds., 20th ed. 1996)).

Pain has been traditionally managed by administering non-opioid analgesics, such as acetylsalicylic acid, choline magnesium trisalicylate, acetaminophen, ibuprofen, fenoprofen, diflusal, and naproxen; or opioid analgesics, including morphine, hydromorphone, methadone, levorphanol, fentanyl, oxycodone, and oxymorphone. *Id.*

UI is uncontrollable urination, generally caused by bladder-detrusor-muscle instability. UI affects people of all ages and levels of physical health, both in health care settings and in the community at large. At present, UI afflicts 15-30% of elderly people living at home, one-third of those living in acute-care settings, and at least one-half of those living in long-term care institutions (R.M. Resnick, *Lancet* 346:94 (1995)). Persons having UI are predisposed to also having urinary-tract infections, pressure ulcers, perineal rashes and urosepsis. Psychosocially, UI is associated with embarrassment, social stigmatization, depression and a risk of institutionalization (Herzo et al., *Annu. Rev. Gerontol. Geriatr.* 9:74 (1989)). Economically, the costs of UI are great; in the United States alone, health-care costs associated with UI are over \$15 billion per annum.

Physiologic bladder contraction results in large part from acetylcholine-induced stimulation of post-ganglionic muscarinic-receptor sites on bladder smooth muscle.

Treatments for UI include the administration of drugs having bladder-relaxant properties, which help to control bladder-detrusor-muscle overactivity. For example, anticholinergics such as propantheline bromide and glycopyrrolate, and combinations of smooth-muscle relaxants such as a combination of racemic oxybutynin and dicyclomine or an
5 anticholinergic, have been used to treat UI (See, e.g., A.J. Wein, *Urol. Clin. N. Am.* 22:557-577 (1995); Levin et al., *J. Urol.* 128:396-398 (1982); Cooke et al., *S. Afr. Med. J.* 63:3 (1983); R.K. Mirakhur et al., *Anaesthesia* 38:1195-1204 (1983)). These drugs are not effective, however, in all patients having uninhibited bladder contractions. Administration of anticholinergic medications represent the mainstay of this type of treatment.

10 None of the existing commercial drug treatments for UI, however, has achieved complete success in all classes of UI patients, nor has treatment occurred without significant adverse side effects. For example, drowsiness, dry mouth, constipation, blurred vision, headaches, tachycardia, and cardiac arrhythmia, which are related to the anticholinergic activity of traditional anti-UI drugs, can occur frequently and adversely affect
15 patient compliance. Yet despite the prevalence of unwanted anticholinergic effects in many patients, anticholinergic drugs are currently prescribed for patients having UI. *The Merck Manual of Medical Information* 631-634 (R. Berkow ed., 1997).

Ulcers are sores occurring where the lining of the digestive tract has been eroded by stomach acids or digestive juices. The sores are typically well-defined round or
20 oval lesions primarily occurring in the stomach and duodenum. About 1 in 10 people develop an ulcer. Ulcers develop as a result of an imbalance between acid-secretory factors, also known as "aggressive factors," such as stomach acid, pepsin, and *Helicobacter pylori* infection, and local mucosal-protective factors, such as secretion of bicarbonate, mucus, and prostaglandins.

25 Treatment of ulcers typically involves reducing or inhibiting the aggressive factors. For example, antacids such as aluminum hydroxide, magnesium hydroxide, sodium bicarbonate, and calcium bicarbonate can be used to neutralize stomach acids. Antacids, however, can cause alkalosis, leading to nausea, headache, and weakness. Antacids can also interfere with the absorption of other drugs into the blood stream and cause diarrhea.

30 H₂ antagonists, such as cimetidine, ranitidine, famotidine, and nizatidine, are also used to treat ulcers. H₂ antagonists promote ulcer healing by reducing gastric acid and digestive-enzyme secretion elicited by histamine and other H₂ agonists in the stomach and duodenum. H₂ antagonists, however, can cause breast enlargement and impotence in men, mental changes (especially in the elderly), headache, dizziness, nausea, myalgia, diarrhea,
35 rash, and fever.

H⁺, K⁺ - ATPase inhibitors such as omeprazole and lansoprazole are also used to treat ulcers. H⁺, K⁺ - ATPase inhibitors inhibit the production of enzymes used by the stomach to secrete acid. Side effects associated with H⁺, K⁺ - ATPase inhibitors include

nausea, diarrhea, abdominal colic, headache, dizziness, somnolence, skin rashes, and transient elevations of plasma activities of aminotransferases.

Sucraflate is also used to treat ulcers. Sucraflate adheres to epithelial cells and is believed to form a protective coating at the base of an ulcer to promote healing. Sucraflate, however, can cause constipation, dry mouth, and interfere with the absorption of other drugs.

Antibiotics are used when *H. pylori* is the underlying cause of the ulcer. Often antibiotic therapy is coupled with the administration of bismuth compounds such as bismuth subsalicylate and colloidal bismuth citrate. The bismuth compounds are believed to enhance secretion of mucous and HCO_3^- , inhibit pepsin activity, and act as an antibacterial against *H. pylori*. Ingestion of bismuth compounds, however, can lead to elevated plasma concentrations of Bi^{+3} and can interfere with the absorption of other drugs.

Prostaglandin analogues, such as misoprostal, inhibit secretion of acid and stimulate the secretion of mucous and bicarbonate and are also used to treat ulcers, especially ulcers in patients who require nonsteroidal anti-inflammatory drugs. Effective oral doses of prostaglandin analogues, however, can cause diarrhea and abdominal cramping. In addition, some prostaglandin analogues are abortifacients.

Carbenoxolone, a mineral corticoid, can also be used to treat ulcers. Carbenoxolone appears to alter the composition and quantity of mucous, thereby enhancing the mucosal barrier. Carbenoxolone, however, can lead to Na^+ and fluid retention, hypertension, hypokalemia, and impaired glucose tolerance.

Muscarinic cholinergic antagonists such as pirenzapine and telenzapine can also be used to reduce acid secretion and treat ulcers. Side effects of muscarinic cholinergic antagonists include dry mouth, blurred vision, and constipation. *The Merck Manual of Medical Information* 496-500 (R. Berkow ed., 1997) and *Goodman and Gilman's The Pharmacological Basis of Therapeutics* 901-915 (J. Hardman and L. Limbird eds., 9th ed. 1996).

IBD is a chronic disorder in which the bowel becomes inflamed, often causing recurring abdominal cramps and diarrhea. The two types of IBD are Crohn's disease and ulcerative colitis.

Crohn's disease, which can include regional enteritis, granulomatous ileitis, and ileocolitis, is a chronic inflammation of the intestinal wall. Crohn's disease occurs equally in both sexes and is more common in Jews of eastern-European ancestry. Most cases of Crohn's disease begin before age 30 and the majority start between the ages of 14 and 24. The disease typically affects the full thickness of the intestinal wall. Generally the disease affects the lowest portion of the small intestine (ileum) and the large intestine, but can occur in any part of the digestive tract.

Early symptoms of Crohn's disease are chronic diarrhea, crampy abdominal pain, fever, loss of appetite, and weight loss. Complications associated with Crohn's disease

include the development of intestinal obstructions, abnormal connecting channels (fistulas), and abscesses. The risk of cancer of the large intestine is increased in people who have Crohn's disease. Often Crohn's disease is associated with other disorders such as gallstones, inadequate absorption of nutrients, amyloidosis, arthritis, episcleritis, aphthous stomatitis, erythema nodosum, pyoderma gangrenosum, ankylosing spondylitis, sacroilitis, uveitis, and primary sclerosing cholangitis. There is no known cure for Crohn's disease.

Cramps and diarrhea, side effects associated with Crohn's disease, can be relieved by anticholinergic drugs, diphenoxylate, loperamide, deodorized opium tincture, or codeine. Generally, the drug is taken orally before a meal.

Broad-spectrum antibiotics are often administered to treat the symptoms of Crohn's disease. The antibiotic metronidazole is often administered when the disease affects the large intestine or causes abscesses and fistulas around the anus. Long-term use of metronidazole, however, can damage nerves, resulting in pins-and-needles sensations in the arms and legs. Sulfasalazine and chemically related drugs can suppress mild inflammation, especially in the large intestine. These drugs, however, are less effective in sudden, severe flare-ups. Corticosteroids, such as prednisone, reduce fever and diarrhea and relieve abdominal pain and tenderness. Long-term corticosteroid therapy, however, invariably results in serious side effects such as high blood-sugar levels, increased risk of infection, osteoporosis, water retention, and fragility of the skin. Drugs such as azathioprine and mercaptourine can compromise the immune system and are often effective for Crohn's disease in patients that do not respond to other drugs. These drugs, however, usually need 3 to 6 months before they produce benefits and can cause serious side effects such as allergy, pancreatitis, and low white-blood-cell count.

When Crohn's disease causes the intestine to be obstructed or when abscesses or fistulas do not heal, surgery can be necessary to remove diseased sections of the intestine. Surgery, however, does not cure the disease, and inflammation tends to recur where the intestine is rejoined. In almost half of the cases a second operation is needed. *The Merck Manual of Medical Information* 528-530 (R. Berkow ed., 1997).

Ulcerative colitis is a chronic disease in which the large intestine becomes inflamed and ulcerated, leading to episodes of bloody diarrhea, abdominal cramps, and fever. Ulcerative colitis usually begins between ages 15 and 30; however, a small group of people have their first attack between ages 50 and 70. Unlike Crohn's disease, ulcerative colitis never affects the small intestine and does not affect the full thickness of the intestine. The disease usually begins in the rectum and the sigmoid colon and eventually spreads partially or completely through out the large intestine. The cause of ulcerative colitis is unknown.

Treatment of ulcerative colitis is directed to controlling inflammation, reducing symptoms, and replacing lost fluids and nutrients. Anticholinergic drugs and low doses of diphenoxylate or loperamide are administered for treating mild diarrhea. For more

intense diarrhea higher doses of diphenoxylate or loperamide, or deodorized opium tincture or codeine are administered. Sulfasalazine, olsalazine, prednisone, or mesalamine can be used to reduce inflammation. Azathioprine and mercaptopurine have been used to maintain remissions in ulcerative-colitis patients who would otherwise need long-term corticosteroid treatment. In severe cases of ulcerative colitis the patient is hospitalized and given corticosteroids intravenously. People with severe rectal bleeding can require transfusions and intravenous fluids. If toxic colitis develops and treatments fail, surgery to remove the large intestine can be necessary. Non-emergency surgery can be performed if cancer is diagnosed, precancerous lesions are detected, or unremitting chronic disease would otherwise make the person an invalid or dependent on high doses of corticosteroids. Complete removal of the large intestine and rectum permanently cures ulcerative colitis. *The Merck Manual of Medical Information* 530-532 (R. Berkow ed., 1997) and *Goodman and Gilman's The Pharmacological Basis of Therapeutics* (J. Hardman and L. Limbird eds., 9th ed. 1996).

IBS is a disorder of motility of the entire gastrointestinal tract, causing abdominal pain, constipation, and/or diarrhea. IBS affects three-times more women than men. In IBS stimuli such as stress, diet, drugs, hormones, or irritants can cause the gastrointestinal tract to contract abnormally. During an episode of IBS contractions of the gastrointestinal tract become stronger and more frequent, resulting in the rapid transit of food and feces through the small intestine, often leading to diarrhea. Cramps result from the strong contractions of the large intestine and increased sensitivity of pain receptors in the large intestine.

There are two major types of IBS. The first type, spastic-colon type, is commonly triggered by eating, and usually produces periodic constipation and diarrhea with pain. Mucous often appears in the stool. The pain can come in bouts of continuous dull aching pain or cramps, usually in the lower abdomen. The person suffering from spastic-colon type IBS can also experience bloating, gas, nausea, headache, fatigue, depression, anxiety, and difficulty concentrating. The second type of IBS usually produces painless diarrhea or constipation. The diarrhea can begin suddenly and with extreme urgency. Often the diarrhea occurs soon after a meal and can sometimes occur immediately upon awakening.

Treatment of IBS typically involves modification of an IBS-patient's diet. Often it is recommended that an IBS patient avoid beans, cabbage, sorbitol, and fructose. A low-fat, high-fiber diet can also help some IBS patients. Regular physical activity can also help keep the gastrointestinal tract functioning properly. Drugs such as propantheline that slow the function of the gastrointestinal tract are generally not effective for treating IBS.

Antidiarrheal drugs, such as diphenoxylate and loperamide, help with diarrhea. *The Merck Manual of Medical Information* 525-526 (R. Berkow ed., 1997).

Many drugs can cause physical and/or psychological addiction. Those most well known types of these drugs include opiates, such as heroin, opium, and morphine;

sympathomimetics, including cocaine and amphetamines; sedative hypnotics, including alcohol, benzodiazepines and barbiturates; and nicotine, which has effects similar to opioids and sympathomimetics. Drug addiction is characterized by a craving or compulsion for taking the drug and an inability to limit its intake. Additionally, drug dependence is associated with drug tolerance, the loss of effect of the drug following repeated administration, and withdrawal, the appearance of physical and behavioral symptoms when the drug is not consumed. Sensitization occurs if repeated administration of a drug leads to an increased response to each dose. Tolerance, sensitization, and withdrawal are phenomena evidencing a change in the central nervous system resulting from continued use of the drug. This change can motivate the addicted individual to continue consuming the drug despite serious social, legal, physical and/or professional consequences. (*See, e.g.*, U.S. Patent No. 6,109,269 to Rise et al.).

Certain pharmaceutical agents have been administered for treating addiction. U.S. Patent No. 5,556,838 to Mayer et al. describes the use of nontoxic NMDA-blocking agents co-administered with an addictive substance to prevent the development of tolerance or withdrawal symptoms. U.S. Patent No. 5,574,052 to Rose et al. describes co-administration of an addictive substance with an antagonist to partially block the pharmacological effects of the substance. U.S. Patent No. 5,075,341 to Mendelson et al. describes the use of a mixed opiate agonist/antagonist to treat cocaine and opiate addiction. U.S. Patent No. 5,232,934 to Downs describes administration of 3 phenoxypyridine to treat addiction. U.S. Patents No. 5,039,680 and 5,198,459 to Imperato et al. describe using a serotonin antagonist to treat chemical addiction. U.S. Patent No. 5,556,837 to Nestler et. al. describes infusing BDNF or NT 4 growth factors to inhibit or reverse neurological adaptive changes that correlate with behavioral changes in an addicted individual. U.S. Patent No. 5,762,925 to Sagan describes implanting encapsulated adrenal medullary cells into an animal's central nervous system to inhibit the development of opioid intolerance. U.S. Patent No. 6,204,284 to Beer et al. describes racemic (\pm)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane for use in the prevention or relief of a withdrawal syndrome resulting from addiction to drugs and for the treatment of chemical dependencies.

Parkinson's disease is a clinical syndrome comprising bradykinesia (slowness and poverty of movement), muscular rigidity, resting tremor (which usually abates during voluntary movement), and an impairment of postural balance leading to disturbance of gait and falling. The features of Parkinson's disease are a loss of pigmented, dopaminergic neurons of the substantia nigra pars compacta and the appearance of intracellular inclusions known as Lewy bodies (*Goodman and Gillman's The Pharmaceutical Basis of Therapeutics* 506 (9th ed. 1996)). Without treatment, Parkinson's disease progresses to a rigid akinetic state in which patients are incapable of caring for themselves. Death frequently results from complications of immobility, including aspiration pneumonia or pulmonary embolism.

Drugs commonly used for the treatment of Parkinson's disease include carbidopa/levodopa, pergolide, bromocriptine, selegiline, amantadine, and trihexyphenidyl hydrochloride. There remains, however, a need for drugs useful for the treatment of Parkinson's disease and having an improved therapeutic profile.

5 Anxiety is a fear, apprehension, or dread of impending danger often accompanied by restlessness, tension, tachycardia, and dyspnea. Other symptoms commonly associated with anxiety include depression, especially accompanied with dysthymic disorder (chronic "neurotic" depression); panic disorder; agoraphobia and other specific phobias; eating disorders; and many personality disorders. Often anxiety is unattached to a clearly
10 identified treatable primary illness. If a primary illness is found, however, it can be desirable to deal with the anxiety at the same time as the primary illness.

 Currently, benzodiazepines are the most commonly used anti-anxiety agents for generalized anxiety disorder. Benzodiazepines, however, carry the risk of producing impairment of cognition and skilled motor functions, particularly in the elderly, which can
15 result in confusion, delirium, and falls with fractures. Sedatives are also commonly prescribed for treating anxiety. The azapirones, such as buspirone, are also used to treat moderate anxiety. The azapirones, however, are less useful for treating severe anxiety accompanied with panic attacks.

 Epilepsy is a disorder characterized by the tendency to have recurring seizures.
20 The etiology commonly consists of lesions in some part of the cortex, such as a tumor; developmental malformation; or damage due to trauma or stroke. In some cases the etiology is genetic. An epileptic seizure can be triggered by repetitive sounds, flashing lights, video games, or touching certain parts of the body. Epilepsy is typically treated with anti-seizure drugs. In epilepsy cases, where anti-seizure drugs are ineffective, and the defect in the brain
25 is isolated to a small area of the brain, surgical removal of that part of the brain can be helpful in alleviating the seizures. In patients who have several sources for the seizures or who have seizures that spread quickly to all parts of the brain, surgical removal of the nerve fibers that connect the two sides of the brain can be helpful.

 Examples of drugs for treating a seizure and epilepsy include carbamazepine,
30 ethosuximide, gabapentin, lamotrigine, phenobarbital, phenytoin, primidone, valproic acid, trimethadione, benzodiazepines, γ -vinyl GABA, acetazolamide, and felbamate. Anti-seizure drugs, however, can have side effects such as drowsiness; hyperactivity; hallucinations; inability to concentrate; central and peripheral nervous system toxicity, such as nystagmus, ataxia, diplopia, and vertigo; gingival hyperplasia; gastrointestinal disturbances such as
35 nausea, vomiting, epigastric pain, and anorexia; endocrine effects such as inhibition of antidiuretic hormone, hyperglycemia, glycosuria, osteomalacia; and hypersensitivity such as scarlatiniform rash, morbilliform rash, Stevens-Johnson syndrome, systemic lupus erythematosus, and hepatic necrosis; and hematological reactions such as red-cell aplasia,

agranulocytosis, thrombocytopenia, aplastic anemia, and megaloblastic anemia. *The Merck Manual of Medical Information* 345-350 (R. Berkow ed., 1997).

5 A seizure is the result of abnormal electrical discharge in the brain. The discharge can involve a small area of the brain and lead to the person only noticing an odd taste or smell or it can involve a large area of the brain and lead to convulsions, *i.e.*, a seizure that causes jerking and spasms of the muscles throughout the body. Convulsions can also result in brief attacks of altered consciousness and loss of consciousness, muscle control, or bladder control. A seizure is often preceded by auras, *i.e.*, unusual sensations of smell, taste, or vision or an intense feeling that a seizure is about to begin. A seizure typically lasts for
10 about 2 to 5 minutes. When the seizure ends the person can have headache, sore muscles, unusual sensations, confusion, and profound fatigue (postictal state). Usually the person cannot remember what happened during the seizure.

A stroke or cerebrovascular accident, is the death of brain tissue (cerebral infarction) resulting from the lack of blood flow and insufficient oxygen to the brain. A
15 stroke can be either ischemic or hemorrhagic. In an ischemic stroke, blood supply to the brain is cut off because of atherosclerosis or a blood clot that has blocked a blood vessel. In a hemorrhagic stroke, a blood vessel bursts preventing normal blood flow and allowing blood to leak into an area of the brain and destroying it. Most strokes develop rapidly and cause brain damage within minutes. In some cases, however, strokes can continue to worsen for
20 several hours or days. Symptoms of strokes vary depending on what part of the brain is effected. Symptoms include loss or abnormal sensations in an arm or leg or one side of the body, weakness or paralysis of an arm or leg or one side of the body, partial loss of vision or hearing, double vision, dizziness, slurred speech, difficulty in thinking of the appropriate word or saying it, inability to recognize parts of the body, unusual movements, loss of
25 bladder control, imbalance, and falling, and fainting. The symptoms can be permanent and can be associated with coma or stupor. Strokes can cause edema or swelling of the brain which can further damage brain tissue. For persons suffering from a stroke, intensive rehabilitation can help overcome the disability caused by impairment of brain tissue. Rehabilitation trains other parts of the brain to assume the tasks previously performed by the
30 damaged part.

Examples of drugs for treating strokes include anticoagulants such as heparin, drugs that break up clots such as streptokinase or tissue plasminogen activator, and drugs that reduce swelling such as mannitol or corticosteroids. *The Merck Manual of Medical Information* 352-355 (R. Berkow ed., 1997).

35 Pruritus is an unpleasant sensation that prompts scratching. Pruritus can be attributed to dry skin, scabies, dermatitis, herpetiformis, atopic dermatitis, pruritus vulvae et ani, miliaria, insect bites, pediculosis, contact dermatitis, drug reactions, urticaria, urticarial eruptions of pregnancy, psoriasis, lichen planus, lichen simplex chronicus, exfoliative

dermatitis, folliculitis, bullous pemphigoid, and fiberglass dermatitis. Conventionally, pruritus is treated by phototherapy with ultraviolet B or PUVA or with therapeutic agents such as naltrexone, nalmefene, danazol, tricyclics, and antidepressants.

5 Selective antagonists of the metabotropic glutamate receptor 5 ("mGluR5") have been shown to exert analgesic activity in in vivo animal models (K. Walker et al., *Neuropharmacology* 40:1-9 (2000) and A. Dogrul et al., *Neuroscience Letters*, 292(2):115-118 (2000)).

10 Selective antagonists of the mGluR5 receptor have also been shown to exert anxiolytic and anti-depressant activity in in vivo animal models (E. Tatarczynska et al., *Br. J. Pharmacol.* 132(7):1423-1430 (2001) and P.J.M. Will et al., *Trends in Pharmacological Sciences* 22(7):331-37 (2001)).

15 Selective antagonists of the mGluR5 receptor have also been shown to exert anti-Parkinson activity in vivo (K. J. Ossowska et al., *Neuropharmacology* 41(4):413-20 (2001) and P.J.M. Will et al., *Trends in Pharmacological Sciences* 22(7):331-37 (2001)).

20 Selective antagonists of the mGluR5 receptor have also been shown to exert anti-dependence activity *in vivo* (C. Chiamulera et al., *Nature Neuroscience* 4(9):873-74 (2001)).


25 International publication no. WO 02/08221 describes diaryl piperazine compounds and related compounds useful as selective modulators of capsaicin receptors and useful in the treatment of chronic and acute pain conditions, itch, and urinary incontinence.

There remains, however, a clear need in the art for new drugs useful for treating or preventing pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression.

Citation of any reference in Section 2 of this application is not to be construed as an admission that such reference is prior art to the present application.

3. SUMMARY OF THE INVENTION

30 The present invention encompasses compounds having the formula (I):



(I)

and pharmaceutically acceptable salts thereof, wherein:

5 R^1 is methyl or halo;

R^3 is:

(a) $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl, or $-(CH_2)_nOR^6$, which is unsubstituted or substituted with one or more of $-CN$, $-OH$, $-halo$, $-NO_2$, $-CH=NR^6$, or $-NR^6OH$; or

10 (b) $-H$, $-halo$, $-NO_2$, $-CN$, or $-NH_2$;

R^4 is $-H$;

R^5 is:

(a) $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, or $-(C_2-C_6)$ alkynyl; or


15 (b) $-(C_3-C_8)$ cycloalkyl, $-(C_5-C_8)$ cycloalkenyl, $-phenyl$, $-naphthyl$, $-(C_{14})aryl$, $-(C_3-C_7)$ heterocycle, $-(C_1-C_6)$ alkylphenyl, $-(C_1-C_6)$ alkylnaphthyl, $-(C_1-C_6)$ alkyl $(C_{14})aryl$, $-(C_1-C_6)$ alkyl (C_3-C_7) heterocycle, $-(C_2-C_6)$ alkenylphenyl, $-(C_2-C_6)$ alkenylnaphthyl, $-(C_2-C_6)$ alkenyl $(C_{14})aryl$, $-(C_2-C_6)$ alkenyl (C_3-C_7) heterocycle, $-(C_2-C_6)$ alkynyl (C_3-C_8) cycloalkyl, $-(C_2-C_6)$ alkynyl (C_5-C_8) cycloalkenyl, $-(C_2-C_6)$ alkynylphenyl, $-(C_2-C_6)$ alkynylnaphthyl, $-(C_2-C_6)$ alkynyl $(C_{14})aryl$, or $-(C_2-C_6)$ alkynyl (C_3-C_7) heterocycle, which is unsubstituted or substituted with one or more R^7 groups;

20 each R^6 is independently $-H$, $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl, $-(C_3-C_8)$ cycloalkyl, $-(C_5-C_8)$ cycloalkenyl, $-phenyl$, $-(C_3-C_5)$ heterocycle, $C(halo)_3$, or $-CH(halo)_2$;

25 each R^7 is independently $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl, $-(CH_2)_n(C_3-C_8)$ cycloalkyl, $-(CH_2)_n(C_5-C_8)$ cycloalkenyl, $-(CH_2)_nphenyl$, $-(CH_2)_nSR^6$, $-(CH_2)_nCH(halo)_2$, $-(CH_2)_nC(halo)_3$, $-halo$, $-(CH_2)_nOR^6$; and

n is an integer ranging from 0 to 2.

The present invention also encompasses compounds having the formula (II):



and pharmaceutically acceptable salts thereof, wherein:

R¹ is methyl or halo;

R³ is:

(a) -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, or -(CH₂)ₙOR⁶, which is unsubstituted or substituted with one or more of -CN, -OH, -halo, -NO₂, -CH=NR⁶, or -NR⁶OH; or

(b) -H, -halo, -NO₂, -CN, or -NH₂;

R⁴ is -H;

R⁵ is:

(a) -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, or -(C₂-C₆)alkynyl; or


(b) -(C₃-C₈)cycloalkyl, -(C₅-C₈)cycloalkenyl, -phenyl, -naphthyl, -(C₁₄)aryl, -(C₃-C₇)heterocycle, -(C₁-C₆)alkylphenyl, -(C₁-C₆)alkylnaphthyl, -(C₁-C₆)alkyl(C₁₄)aryl, -(C₁-C₆)alkyl(C₃-C₇)heterocycle, -(C₂-C₆)alkenylphenyl, -(C₂-C₆)alkenylphenyl, -(C₂-C₆)alkenyl(C₁₄)aryl, -(C₂-C₆)alkenyl(C₃-C₇)heterocycle, -(C₂-C₆)alkynyl(C₃-C₈)cycloalkyl, -(C₂-C₆)alkynyl(C₅-C₈)cycloalkenyl, -(C₂-C₆)alkynylphenyl, -(C₂-C₆)alkylnaphthyl, -(C₂-C₆)alkynyl(C₁₄)aryl, or -(C₂-C₆)alkynyl(C₃-C₇)heterocycle, which is unsubstituted or substituted with one or more R⁷ groups;

each R⁶ is independently -H, -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -(C₃-C₈)cycloalkyl, -(C₅-C₈)cycloalkenyl, -phenyl, -(C₃-C₅)heterocycle, C(halo)₃, or CH(halo)₂;

each R^7 is independently $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, $-(C_2-C_6)alkynyl$, $-(CH_2)_n(C_3-C_8)cycloalkyl$, $-(CH_2)_n(C_5-C_8)cycloalkenyl$, $-(CH_2)_nphenyl$, $-(CH_2)_nSR^6$, $-(CH_2)_nCH(halo)_2$, $-(CH_2)_nC(halo)_3$, $-halo$, $-(CH_2)_nOR^6$; and

n is an integer ranging from 0 to 2.

The present invention also encompasses compounds having the formula (III):



and pharmaceutically acceptable salts thereof, wherein:

X is O or S;

R^1 is $-CH_3$ or $-halo$;

R^3 is:

(a) $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, $-(C_2-C_6)alkynyl$, or $-(CH_2)_nOR^6$, which is unsubstituted or substituted with one or more of $-CN$, $-OH$, $-halo$, $-NO_2$, $-CH=NR^6$, or $-NR^6OH$; or

(b) $-H$, $-halo$, $-NO_2$, $-CN$, or $-NH_2$;

R^4 is $-H$;

each R^8 is independently $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, $-(C_2-C_6)alkynyl$, $-(CH_2)_n(C_3-C_8)cycloalkyl$, $-(CH_2)_n(C_5-C_8)cycloalkenyl$, $-(CH_2)_nphenyl$, $-(CH_2)_nSR^6$, $-(CH_2)_nCH(halo)_2$, $-(CH_2)_nC(halo)_3$, $-halo$, $-(CH_2)_nOR^6$;

each R^6 is independently $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, $-(C_2-C_6)alkynyl$, $phenyl$, $-naphthyl$, or $-(C_{14})aryl$;


Ar is -phenyl, -naphthyl, $-(C_{14})$ aryl, $-(C_3-C_8)$ cycloalkyl or $-(C_5-C_8)$ cycloalkenyl, which is unsubstituted or substituted with one or more R^7 groups;

each R^7 is independently $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl, $-(CH_2)_n(C_3-C_8)$ cycloalkyl, $-(CH_2)_n(C_5-C_8)$ cycloalkenyl, $-(CH_2)_n$ phenyl, $-(CH_2)_nSR^6$,
 5 $-(CH_2)_nCH(halo)_2$, $-(CH_2)_nC(halo)_3$, -halo, $-(CH_2)_nOR^6$;

n is an integer ranging from 0 to 2; and

p is an integer ranging from 0 to 2.

The present invention also encompasses compounds having the formula (IV):



and pharmaceutically acceptable salts thereof, wherein:

X is O or S;

25 R^1 is $-CH_3$ or -halo;

R^3 is:

(a) $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl, or $-(CH_2)_nOR^6$, which is unsubstituted or substituted with one or more of -CN, -OH, -halo, - NO_2 , $-CH=NR^6$, or $-NR^6OH$; or

30 (b) -H, -halo, $-NO_2$, -CN, or $-NH_2$;

R^4 is -H;

each R^8 is independently $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl,

$-(CH_2)_n(C_3-C_8)\text{cycloalkyl}$, $-(CH_2)_n(C_5-C_8)\text{cycloalkenyl}$, $-(CH_2)_n\text{phenyl}$, $-(CH_2)_nSR^6$,
 $-(CH_2)_nCH(\text{halo})_2$, $-(CH_2)_nC(\text{halo})_3$, -halo, $-(CH_2)_nOR^6$;

each R^6 is independently $-(C_1-C_6)\text{alkyl}$, $-(C_2-C_6)\text{alkenyl}$, $-(C_2-C_6)\text{alkynyl}$, -phenyl, -naphthyl, or $-(C_{14})\text{aryl}$;

5 Ar is -phenyl, -naphthyl, $-(C_{14})\text{aryl}$, $-(C_3-C_8)\text{cycloalkyl}$ or $-(C_5-C_8)\text{cycloalkenyl}$, which is unsubstituted or substituted with one or more R^7 groups;

each R^7 is independently $-(C_1-C_6)\text{alkyl}$, $-(C_2-C_6)\text{alkenyl}$, $-(C_2-C_6)\text{alkynyl}$,
 $-(CH_2)_n(C_3-C_8)\text{cycloalkyl}$, $-(CH_2)_n(C_5-C_8)\text{cycloalkenyl}$, $-(CH_2)_n\text{phenyl}$, $-(CH_2)_nSR^6$,
 $-(CH_2)_nCH(\text{halo})_2$, $-(CH_2)_nC(\text{halo})_3$, -halo, $-(CH_2)_nOR^6$;

10 n is an integer ranging from 0 to 2; and

p is an integer ranging from 0 to 2.

A compound of formula (I) - (IV) or a pharmaceutically acceptable salt thereof (a "Thiadiazolylpiperazine Compound") is useful for treating or preventing pain, UI, an ulcer,
 15 IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression in an animal.

The invention also relates to compositions comprising an effective amount of
 20 a Thiadiazolylpiperazine Compound and a pharmaceutically acceptable carrier or excipient. The compositions are useful for treating or preventing pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting,
 25 dyskinesia, or depression in an animal.

The invention further relates to methods for treating pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a
 30 migraine, vomiting, dyskinesia, or depression comprising administering to an animal in need thereof an effective amount of a Thiadiazolylpiperazine Compound.

The invention further relates to methods for preventing pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted
 35 brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a

migraine, vomiting, dyskinesia, or depression comprising administering to an animal in need thereof an effective amount of a Thiadiazolylpiperazine Compound.

The invention still further relates to methods for inhibiting Vanilloid Receptor 1 ("VR1") function in a cell, comprising contacting a cell capable of expressing VR1 with an effective amount of a Thiadiazolylpiperazine Compound.

The invention still further relates to methods for inhibiting mGluR5 function in a cell, comprising contacting a cell capable of expressing mGluR5 with an effective amount of a Thiadiazolylpiperazine Compound.

The invention still further relates to methods for inhibiting metabotropic glutamate receptor 1 ("mGluR1") function in a cell, comprising contacting a cell capable of expressing mGluR1 with an effective amount of a Thiadiazolylpiperazine Compound.

The invention still further relates to methods for preparing a composition comprising the step of admixing a Thiadiazolylpiperazine Compound and a pharmaceutically acceptable carrier or excipient.


The invention still further relates to a kit comprising a container containing an effective amount of a Thiadiazolylpiperazine Compound.

The present invention can be understood more fully by reference to the following detailed description and illustrative examples, which are intended to exemplify non-limiting embodiments of the invention.

4. DETAILED DESCRIPTION OF THE INVENTION

4.1 THIADIAZOLYPIPERAZINE COMPOUNDS OF FORMULA (I)

As stated above, the present invention encompasses compounds of Formula (I)



(I)

and pharmaceutically acceptable salts thereof, where R^1 , R^3 , R^4 , and R^5 are defined above.

In one embodiment, R^1 is methyl.

In another embodiment, R^1 is halo.

In another embodiment, R^3 is $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, $-(C_2-C_6)alkynyl$, or $-(CH_2)_nOR^6$, which is unsubstituted or substituted with one or more of $-CN$, $-OH$, $-halo$, $-NO_2$, $-CH=NR^6$, or $-NR^6OH$.

In another embodiment, R³ is -H, -halo, -NO₂, -CN, or -NH₂.

In another embodiment, R³ is -H;

In another embodiment, R³ is -CH₃.

In another embodiment, R⁵ is -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, or -(C₂-
5 C₆)alkynyl.

In another embodiment, R⁵ is -(C₃-C₈)cycloalkyl, -(C₅-C₈)cycloalkenyl, -
phenyl, -naphthyl, -(C₁₄)aryl, -(C₃-C₇)heterocycle, -(C₁-C₆)alkylphenyl, -(C₁-C₆)alkylnaphthyl, -
(C₁-C₆)alkyl(C₁₄)aryl, -(C₁-C₆)alkyl(C₃-C₇)heterocycle, -(C₂-C₆)alkenylphenyl, -(C₂-
C₆)alkenylnaphthyl, -(C₂-C₆)alkenyl(C₁₄)aryl, -(C₂-C₆)alkenyl(C₃-C₇)heterocycle, -(C₂-
10 C₆)alkynyl(C₃-C₈)cycloalkyl, -(C₂-C₆)alkynyl(C₅-C₈)cycloalkenyl, -(C₂-C₆)alkynylphenyl, -
(C₂-C₆)alkenylnaphthyl, -(C₂-C₆)alkynyl(C₁₄)aryl, or -(C₂-C₆)alkynyl(C₃-C₇)heterocycle,
which is unsubstituted or substituted with one or more R⁷ groups.

In another embodiment, R⁵ is phenyl optionally substituted with a -(C₁-
C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is substituted at the phenyl
15 group's 4-position. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.


In another embodiment, R¹ is methyl and R⁵ is phenyl. In another
embodiment, R¹ is methyl and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group. In another
embodiment, R¹ is methyl and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group at the
phenyl group's 4-position. In another embodiment, R¹ methyl and R⁵ is phenyl substituted
20 with a *t*-butyl group at the phenyl group's 4-position.

In another embodiment, R¹ is -halo and R⁵ is phenyl. In another embodiment,
R¹ is -halo and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group. In another embodiment,
R¹ -halo and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group at the phenyl group's 4-
position. In another embodiment, R¹ -halo and R⁵ is phenyl substituted with a *t*-butyl group
25 at the phenyl group's 4-position.

In another embodiment, R¹ is -chloro and R⁵ is phenyl. In another
embodiment, R¹ is -chloro and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group. In
another embodiment, R¹ -chloro and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group at
the phenyl group's 4-position. In another embodiment, R¹ -chloro and R⁵ is phenyl
30 substituted with a *t*-butyl group at the phenyl group's 4-position.

4.2 THIADIAZOLYLPIPERAZINE COMPOUNDS OF FORMULA (II)

The present invention also encompasses compounds of Formula (II)



and pharmaceutically acceptable salts thereof, where R^1 , R^3 , R^4 , and R^5 are defined above.

5 In one embodiment, R^1 is methyl.

In another embodiment, R^1 is halo.

In another embodiment, R^3 is $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, $-(C_2-C_6)$ alkynyl, or $-(CH_2)_nOR^6$, which is unsubstituted or substituted with one or more of $-CN$, $-OH$, $-halo$, $-NO_2$, $-CH=NR^6$, or $-NR^6OH$.

10 In another embodiment, R^3 is $-H$, $-halo$, $-NO_2$, $-CN$, or $-NH_2$.

In another embodiment, R^3 is $-H$;

In another embodiment, R^3 is $-CH_3$.

In another embodiment, R^5 is $-(C_1-C_6)$ alkyl, $-(C_2-C_6)$ alkenyl, or $-(C_2-C_6)$ alkynyl.

15 In another embodiment, R^5 is $-(C_3-C_8)$ cycloalkyl, $-(C_5-C_8)$ cycloalkenyl, $-phenyl$, $-naphthyl$, $-(C_{14})aryl$, $-(C_3-C_7)$ heterocycle, $-(C_1-C_6)$ alkylphenyl, $-(C_1-C_6)$ alkylnaphthyl, $-(C_1-C_6)$ alkyl $(C_{14})aryl$, $-(C_1-C_6)$ alkyl (C_3-C_7) heterocycle, $-(C_2-C_6)$ alkenylphenyl, $-(C_2-C_6)$ alkenylnaphthyl, $-(C_2-C_6)$ alkenyl $(C_{14})aryl$, $-(C_2-C_6)$ alkenyl (C_3-C_7) heterocycle, $-(C_2-C_6)$ alkynyl (C_3-C_8) cycloalkyl, $-(C_2-C_6)$ alkynyl (C_5-C_8) cycloalkenyl, $-(C_2-C_6)$ alkynylphenyl, $-(C_2-C_6)$ alkynylnaphthyl, $-(C_2-C_6)$ alkynyl $(C_{14})aryl$, or $-(C_2-C_6)$ alkynyl (C_3-C_7) heterocycle, which is unsubstituted or substituted with one or more R^7 groups.

In another embodiment, R^5 is phenyl optionally substituted with a $-(C_1-C_6)$ alkyl group. In another embodiment, the $-(C_1-C_6)$ alkyl group is substituted at the phenyl group's 4-position. In another embodiment, the $-(C_1-C_6)$ alkyl group is a *t*-butyl group.


25 In another embodiment, R^1 is methyl and R^5 is phenyl. In another embodiment, R^1 is methyl and R^5 is phenyl substituted with a $-(C_1-C_6)$ alkyl group. In another embodiment, R^1 is methyl and R^5 is phenyl substituted with a $-(C_1-C_6)$ alkyl group at the phenyl group's 4-position. In another embodiment, R^1 methyl and R^5 is phenyl substituted with a *t*-butyl group at the phenyl group's 4-position.

30 In another embodiment, R^1 is $-halo$ and R^5 is phenyl. In another embodiment, R^1 is $-halo$ and R^5 is phenyl substituted with a $-(C_1-C_6)$ alkyl group. In another embodiment, R^1 $-halo$ and R^5 is phenyl substituted with a $-(C_1-C_6)$ alkyl group at the phenyl group's 4-position. In another embodiment, R^1 $-halo$ and R^5 is phenyl substituted with a *t*-butyl group at the phenyl group's 4-position.

In another embodiment, R¹ is -chloro and R⁵ is phenyl. In another embodiment, R¹ is -chloro and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group. In another embodiment, R¹ -chloro and R⁵ is phenyl substituted with a -(C₁-C₆)alkyl group at the phenyl group's 4-position. In another embodiment, R¹ -chloro and R⁵ is phenyl substituted with a *t*-butyl group at the phenyl group's 4-position.

4.3 THIADIAZOLYLPIPERAZINE COMPOUNDS OF FORMULA (III)

The present invention also encompasses compounds of Formula (III)



(III)

and pharmaceutically acceptable salts thereof, where R¹, R³, R⁴, R⁸, X, Ar, and p are defined above.

In one embodiment, X is O.

In another embodiment, X is S.

In another embodiment, R¹ is methyl.

In another embodiment, R¹ is -halo.

In another embodiment R¹ is -chloro.

In another embodiment, R¹ is -bromo.

In another embodiment, R¹ is -iodo.

In another embodiment, R¹ is -fluoro.

In another embodiment, X is O and R¹ is methyl.

In another embodiment, X is O and R¹ is -halo.

In another embodiment X is O and R¹ is -chloro.

In another embodiment, X is O and R¹ is -bromo.

In another embodiment, X is O and R¹ is -iodo.

In another embodiment, X is O and R¹ is -fluoro.

In another embodiment, X is S and R¹ is methyl.

In another embodiment, X is S and R¹ is -halo.

In another embodiment X is S and R¹ is -chloro.

In another embodiment, X is S and R¹ is -bromo.

In another embodiment, X is S and R¹ is -iodo.

5 In another embodiment, X is S and R¹ is -fluoro.

In another embodiment, R³ is -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, or -(CH₂)_nOR⁶, which is unsubstituted or substituted with one or more of -CN, -OH, -halo, -NO₂, -CH=NR⁶, or -NR⁶OH.

In another embodiment, R³ is -H, -halo, -NO₂, -CN, or -NH₂.

10 In another embodiment, R³ is -H.

In another embodiment, R³ is -CH₃.

In another embodiment, Ar is -phenyl, which is unsubstituted or substituted with one or more R⁷ groups.

15 In another embodiment, Ar is -naphthyl, which is unsubstituted or substituted with one or more R⁷ groups.

In another embodiment, Ar is -(C₁₄)aryl, which is unsubstituted or substituted with one or more R⁷ groups.

In another embodiment, Ar is -(C₃-C₈)cycloalkyl, which is unsubstituted or substituted with one or more R⁷ groups.

20 In another embodiment, Ar is -(C₅-C₈)cycloalkenyl, which is unsubstituted or substituted with one or more R⁷ groups.

In another embodiment, R¹ is methyl and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group.

25 In another embodiment, R¹ is methyl and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

In another embodiment, R¹ is -halo and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group.

30 In another embodiment, R¹ is -halo and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

In another embodiment, R¹ is -chloro and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group

35 In another embodiment, R¹ is -chloro and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

In another embodiment, R¹ is -bromo and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group

In another embodiment, R^1 is -bromo and Ar is phenyl, substituted in the 4-position with a $-(C_1-C_6)$ alkyl group. In another embodiment, the $-(C_1-C_6)$ alkyl group is a *t*-butyl group.

In another embodiment, R^1 is -iodo and Ar is phenyl, optionally substituted with a $-(C_1-C_6)$ alkyl group

In another embodiment, R^1 is -iodo and Ar is phenyl, substituted in the 4-position with a $-(C_1-C_6)$ alkyl group. In another embodiment, the $-(C_1-C_6)$ alkyl group is a *t*-butyl group.

In another embodiment, R^1 is -fluoro and Ar is phenyl, optionally substituted with a $-(C_1-C_6)$ alkyl group

In another embodiment, R^1 is -fluoro and Ar is phenyl, substituted in the 4-position with a $-(C_1-C_6)$ alkyl group. In another embodiment, the $-(C_1-C_6)$ alkyl group is a *t*-butyl group.

In another embodiment, p is 0.

In another embodiment, p is 1.


In another embodiment, p is 2.

In another embodiment, p is 1 and R^8 is $-CH_3$.

In another embodiment, p is 2 and each R^8 is $-CH_3$.

4.4 THIADIAZOLYLPIPERAZINE COMPOUNDS OF FORMULA (IV)

The present invention also encompasses compounds of Formula (IV)



and pharmaceutically acceptable salts thereof, where R^1 , R^3 , R^4 , R^8 , X , Ar , and p are defined above.

In one embodiment, X is O .

In another embodiment, X is S .

- In another embodiment, R¹ is methyl.
 In another embodiment, R¹ is -halo.
 In another embodiment R¹ is -chloro.
 In another embodiment, R¹ is -bromo.
 5 In another embodiment, R¹ is -iodo.
 In another embodiment, R¹ is -fluoro.
 In another embodiment, X is O and R¹ is methyl.
 In another embodiment, X is O and R¹ is -halo.
 In another embodiment X is O and R¹ is -chloro.
 10 In another embodiment, X is O and R¹ is -bromo.
 In another embodiment, X is O and R¹ is -iodo.
 In another embodiment, X is O and R¹ is -fluoro.
 In another embodiment, X is S and R¹ is methyl.
 In another embodiment, X is S and R¹ is -halo.
 15 In another embodiment X is S and R¹ is -chloro.
 In another embodiment, X is S and R¹ is -bromo.
 In another embodiment, X is S and R¹ is -iodo.
 In another embodiment, X is S and R¹ is -fluoro.
 In another embodiment, R³ is -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl,
 20 or -(CH₂)_nOR⁶, which is unsubstituted or substituted with one or more of -CN, -OH, -halo, -
 NO₂, -CH=NR⁶, or -NR⁶OH.
 In another embodiment, R³ is -H, -halo, -NO₂, -CN, or -NH₂.
 In another embodiment, R³ is -H.
 In another embodiment, R³ is -CH₃.
 25 In another embodiment, Ar is -phenyl, which is unsubstituted or substituted
 with one or more R⁷ groups.
 In another embodiment, Ar is -naphthyl, which is unsubstituted or substituted
 with one or more R⁷ groups.
 In another embodiment, Ar is -(C₁₄)aryl, which is unsubstituted or substituted
 30 with one or more R⁷ groups.
 In another embodiment, Ar is -(C₃-C₈)cycloalkyl, which is unsubstituted or
 substituted with one or more R⁷ groups.
 In another embodiment, Ar is -(C₅-C₈)cycloalkenyl, which is unsubstituted or
 substituted with one or more R⁷ groups.
 35 In another embodiment, R¹ is methyl and Ar is phenyl, optionally substituted
 with a -(C₁-C₆)alkyl group.

In another embodiment, R¹ is methyl and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

5 In another embodiment, R¹ is -halo and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group.

In another embodiment, R¹ is -halo and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

10 In another embodiment, R¹ is -chloro and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group.

In another embodiment, R¹ is -chloro and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

15 In another embodiment, R¹ is -bromo and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group.

In another embodiment, R¹ is -bromo and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

20 In another embodiment, R¹ is -iodo and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group.

In another embodiment, R¹ is -iodo and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

25 In another embodiment, R¹ is -fluoro and Ar is phenyl, optionally substituted with a -(C₁-C₆)alkyl group

In another embodiment, R¹ is -fluoro and Ar is phenyl, substituted in the 4-position with a -(C₁-C₆)alkyl group. In another embodiment, the -(C₁-C₆)alkyl group is a *t*-butyl group.

30 In another embodiment, p is 0.

In another embodiment, p is 1.

In another embodiment, p is 2

In another embodiment, p is 1 and R⁸ is -CH₃.


In another embodiment, p is 2 and each R⁸ is -CH₃.

35 In another embodiment, a composition or pharmaceutical composition is prepared by admixing a Thiadiazolylpiperazine Compound or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier or excipient. Admixing can be accomplished using methods well known for admixing a compound (or a salt) and a pharmaceutically acceptable carrier or excipient. In another embodiment, the

Thiadiazolylpiperazine Compound or a pharmaceutically acceptable salt thereof is present in an effective amount.

Illustrative Thiadiazolylpiperazine Compounds, obtainable by the methods described herein, are listed below in Tables (I) and (II):

Table (I)



and pharmaceutically acceptable salts thereof, wherein

Compound	R ¹	R ³	R ⁵
AAA	-CH ₃	-H	-phenyl
AAB	-CH ₃	-H	-(4-(<i>isopropyl</i>)phenyl
AAC	-CH ₃	-H	-(4-trifluoromethyl)phenyl
AAD	-CH ₃	-H	-(4-bromo)phenyl
AAE	-CH ₃	-H	-(4-chloro)phenyl
AAF	-CH ₃	-H	-(4-methyl)phenyl
AAG	-CH ₃	-H	-(4- <i>tert</i> butyl)phenyl
AAH	-CH ₃	-H	-2-(cyclopropyl)phenyl
AAI	-CH ₃	-H	-2(4-bromophenyl)ethyl
AAJ	-CH ₃	-H	-2(4-methylphenyl)ethyl
AAK	-CH ₃	-CH ₃	-phenyl
AAL	-CH ₃	-CH ₃	-(4-(<i>isopropyl</i>)phenyl
AAM	-CH ₃	-CH ₃	-(4-trifluoromethyl)phenyl
AAN	-CH ₃	-CH ₃	-(4-bromo)phenyl
AAO	-CH ₃	-CH ₃	-(4-chloro)phenyl
AAP	-CH ₃	-CH ₃	-(4-methyl)phenyl
AAQ	-CH ₃	-CH ₃	-(4- <i>tert</i> butyl)phenyl
AAR	-CH ₃	-CH ₃	-2-(cyclopropyl)phenyl
AAS	-CH ₃	-CH ₃	-2(4-bromophenyl)ethyl


Compound	R ¹	R ³	R ⁵
AAT	-CH ₃	-CH ₃	-2(4-methylphenyl)ethyl
AAU	-Cl	-H	-phenyl
AAV	-Cl	-H	-(4-(<i>isopropyl</i>)phenyl)
AAW	-Cl	-H	-(4-trifluoromethyl)phenyl
AAX	-Cl	-H	-(4-bromo)phenyl
AAZ	-Cl	-H	-(4-chloro)phenyl
AAZ	-Cl	-H	-(4-methyl)phenyl
ABA	-Cl	-H	-(4- <i>tert</i> butyl)phenyl
ABB	-Cl	-H	-2-(cyclopropyl)phenyl
ABC	-Cl	-H	-2(4-bromophenyl)ethyl
ABD	-Cl	-H	-2(4-methylphenyl)ethyl
ABE	-Cl	-CH ₃	-phenyl
ABF	-Cl	-CH ₃	-(4-(<i>isopropyl</i>)phenyl)
ABG	-Cl	-CH ₃	-(4-trifluoromethyl)phenyl
ABH	-Cl	-CH ₃	-(4-bromo)phenyl
ABI	-Cl	-CH ₃	-(4-chloro)phenyl
ABJ	-Cl	-CH ₃	-(4-methyl)phenyl
ABK	-Cl	-CH ₃	-(4- <i>tert</i> butyl)phenyl
ABL	-Cl	-CH ₃	-2-(cyclopropyl)phenyl
ABM	-Cl	-CH ₃	-2(4-bromophenyl)ethyl
ABN	-Cl	-CH ₃	-2(4-methylphenyl)ethyl

5

10

15

Table (II)



and pharmaceutically acceptable salts thereof, wherein

Compound	R ¹	R ³	R ⁵
ABO	-CH ₃	-H	-phenyl
ABP	-CH ₃	-H	-(4-(<i>isopropyl</i>)phenyl
ABQ	-CH ₃	-H	-(4-(trifluoromethyl)phenyl
ABR	-CH ₃	-H	-(4-bromo)phenyl
ABS	-CH ₃	-H	-(4-chloro)phenyl
ABT	-CH ₃	-H	-(4-methyl)phenyl
ABU	-CH ₃	-H	-(4- <i>tert</i> butyl)phenyl
ABV	-CH ₃	-H	-2-(cyclopropyl)phenyl
ABW	-CH ₃	-H	-2(4-bromophenyl)ethyl
ABX	-CH ₃	-H	-2(4-methylphenyl)ethyl
ABY	-CH ₃	-CH ₃	-phenyl
ABZ	-CH ₃	-CH ₃	-(4-(<i>isopropyl</i>)phenyl
ACA	-CH ₃	-CH ₃	-(4-(trifluoromethyl)phenyl
ACB	-CH ₃	-CH ₃	-(4-bromo)phenyl
ACC	-CH ₃	-CH ₃	-(4-chloro)phenyl
ACD	-CH ₃	-CH ₃	-(4-methyl)phenyl
ACE	-CH ₃	-CH ₃	-(4- <i>tert</i> butyl)phenyl
ACF	-CH ₃	-CH ₃	-2-(cyclopropyl)phenyl
ACG	-CH ₃	-CH ₃	-2(4-bromophenyl)ethyl
ACH	-CH ₃	-CH ₃	-2(4-methylphenyl)ethyl
ACI	-Cl	-H	-phenyl
ACJ	-Cl	-H	-(4-(<i>isopropyl</i>)phenyl
ACK	-Cl	-H	-(4-(trifluoromethyl)phenyl

ACL	-Cl	-H	-(4-bromo)phenyl
ACM	-Cl	-H	-(4-chloro)phenyl
CAN	-Cl	-H	-(4-methyl)phenyl
ACO	-Cl	-H	-(4- <i>tert</i> butyl)phenyl
ACP	-Cl	-H	-2-(cyclopropyl)phenyl
ACQ	-Cl	-H	-2(4-bromophenyl)ethyl
ACR	-Cl	-H	-2(4-methylphenyl)ethyl
ACS	-Cl	-CH ₃	-phenyl
ACT	-Cl	-CH ₃	-(4-(<i>isopropyl</i>)phenyl
ACU	-Cl	-CH ₃	-(4-(trifluoromethyl)phenyl
ACV	-Cl	-CH ₃	-(4-bromo)phenyl
ACW	-Cl	-CH ₃	-(4-chloro)phenyl
ACX	-Cl	-CH ₃	-(4-methyl)phenyl
ACY	-Cl	-CH ₃	-(4- <i>tert</i> butyl)phenyl
ACZ	-Cl	-CH ₃	-2-(cyclopropyl)phenyl
ADA	-Cl	-CH ₃	-2(4-bromophenyl)ethyl
ADB	-Cl	-CH ₃	-2(4-methylphenyl)ethyl

As used herein, in connection with the Thiadiazolylpiperazine compounds, the terms used above having following meaning:

5 “-Alkyl” means a straight chain or branched non-cyclic hydrocarbon having from 1 to 10 carbon atoms. “-(C₁-C₄)alkyl” means an alkyl group having from 1 and 4 carbon atoms. “-(C₁-C₆)alkyl” means an alkyl group having from 1 and 6 carbon atoms. “-(C₂-C₆)alkyl” means an alkyl group having from 2 and 6 carbon atoms. Representative straight chain alkyls include -methyl, -ethyl, -n-propyl, -n-butyl, -n-pentyl, -n-hexyl, -n-heptyl, -n-octyl, -n-nonyl and -n-decyl; while branched alkyls include -isopropyl, -*sec*-butyl, -*tert*-butyl, -isopentyl, 2-methylbutyl, and the like.

10 “-Alkenyl” means a straight chain or branched non-cyclic hydrocarbon having from 1 to 10 carbon atoms and including at least one carbon-carbon double bond. “-(C₂-C₆)alkenyl” means an alkenyl group having from 2 and 6 carbon atoms. “-(C₂-C₃)alkenyl” means an alkenyl group having from 2 and 3 carbon atoms. Representative straight chain and branched alkenyls include -vinyl, -allyl, -1-butenyl, -2-butenyl, -isobutenyl, -1-pentenyl, -2-pentenyl, -3-methyl-1-butenyl, -2-methyl-2-butenyl, -2,3-dimethyl-2-butenyl, 1-hexyl, 2-hexyl, 3-hexyl, and the like.

“-Alkynyl” means a straight chain or branched non-cyclic hydrocarbon having from 1 to 10 carbon atoms and including at least one carbon-carbon triple bond. “-(C₂-C₆)alkynyl” means an alkynyl group having from 2 and 6 carbon atoms. Representative straight chain and branched alkynyls include -acetylenyl, -propynyl, -1-butylnyl, -2-butylnyl, -1-pentylnyl, -2-pentylnyl, -3-methyl-1 butynyl, and the like.

“-Cycloalkyl” means a saturated cyclic hydrocarbon having from 3 to 10 carbon atoms. “-(C₃-C₈)cycloalkyl” means a cycloalkyl group having from 3 and 8 carbon atoms. Representative cycloalkyls include -cyclopropyl, -cyclobutyl, -cyclopentyl, -cyclohexyl, -cycloheptyl, -cyclooctyl, -cyclononyl, and -cyclodecyl. Cycloalkyls also include bi- and tri-cyclic ring systems having from 8 to 14 carbon atoms such as a cycloalkyl (such as cyclopentane or cyclohexane) fused to one or more aromatic (such as phenyl) or non-aromatic (such as cyclohexane) carbocyclic rings.

“-Cycloalkenyl” means a cyclic hydrocarbon having at least one carbon-carbon double bond in the cyclic system and from 5 to 10 carbon atoms. “-(C₅-C₈)cycloalkenyl” means a cycloalkenyl group having from 5 and 8 carbon atoms. Representative cycloalkenyl include -cyclopentenyl, -cyclopentadienyl, -cyclohexenyl, -cyclohexadienyl, cycloheptenyl, cycloheptadienyl, cycloheptatrienyl, cyclooctenyl, cyclooctadienyl, cyclooctatrienyl, cyclooctatetraenyl, and the like. Cycloalkenyls also include bi- and tri-cyclic ring systems having from 8 to 14 carbon atoms such as a cycloalkenyl (such as cyclopentene or cyclohexene) fused to one or more aromatic (such as phenyl) or non-aromatic (such as cyclohexane) carbocyclic rings.

“-Halogen” or “-Halo” means fluorine, chlorine, bromine or iodine.

“-Aryl” means an aromatic carbocyclic moiety such as phenyl, naphthyl and anthracenyl. “-(C₁₄)aryl” means an aryl group having 14 carbon atoms.

“-Heteroaryl” means an aromatic heterocycle ring of 5 to 10 members and having at least one heteroatom selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom, including both mono- and bicyclic ring systems. Representative heteroaryls are pyridyl, furyl, benzofuranyl, thiophenyl, benzothiophenyl, quinolinyl, pyrrolyl, indolyl, oxazolyl, benzoxazolyl, imidazolyl, benzimidazolyl, thiazolyl, benzothiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, cinnolinyl, phthalazinyl, and quinazolinyl.

“-Heterocycle” or “-Heterocyclo” means a 3- to 7-membered monocyclic, or 7- to 10-membered bicyclic, heterocyclic ring which is either saturated, unsaturated, or aromatic, and which contains from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulfur, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen heteroatom may be optionally quaternized, including bicyclic rings in which any of the above heterocycles are fused to a benzene ring. The heterocycle may be attached via any heteroatom or carbon atom. Heterocycles include heteroaryls as defined

above. “-(C₃-C₅)heterocycle” means a heterocycle having from 3 and 5 ring atoms, 1 to 4 of which are heteroatoms. “-(C₃-C₇)heterocycle” means a heterocycle having from 3 and 7 ring atoms, 1 to 4 of which are heteroatoms. Thus, in addition to the heteroaryls listed above, heterocycles also include morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyrindinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like.

“-Alkylaryl” means an alkyl having at least one alkyl hydrogen atom replaced with an aryl moiety, defined above. Illustrative -alkylaryl groups include benzyl,

-(CH₂)₂phenyl, -(CH₂)₃phenyl, -CH(phenyl)₂, and the like.

“-Alkylheterocycle” means an alkyl group wherein at least one hydrogen of the alkyl group is substituted with a heterocycle group, defined above.

“-Alkenylaryl” means an alkenyl having at least one hydrogen atom of the alkenyl group replaced with an aryl group, defined above.

“-Alkenylheterocycle” means an alkenyl having at least one hydrogen atom of the alkenyl group replaced with a heterocycle group, defined above.

“-Alkynylcycloalkyl” means an alkynyl having at least one hydrogen atom of the alkynyl group replaced with a cycloalkyl group, defined above.

“-Alkynylcycloalkenyl” means an alkynyl having at least one hydrogen atom of the alkynyl group replaced with a cycloalkenyl group, defined above.

“-Alkynylaryl” means an alkynyl having at least one hydrogen atom of the alkynyl group replaced with an aryl group, defined above.

“-Alkynylheterocycle” means an alkynyl having at least one hydrogen atom of the alkynyl group replaced with a heterocycle group, defined above.

The Thiadiazolylpiperazine Compounds can have a substituent, for example R₅, having a first and a second group. Examples of substituents having a first and a second group are -(C₁-C₆)alkylphenyl, -(C₁-C₆)alkylnaphthyl, -(C₁-C₆)alkyl(C₁₄)aryl, -(C₁-C₆)alkyl(C₃-C₇)heterocycle, -(C₂-C₆)alkenylphenyl, -(C₂-C₆)alkenylnaphthyl, -(C₂-C₆)alkenyl(C₁₄)aryl, -(C₂-C₆)alkenyl(C₃-C₇)heterocycle, -(C₂-C₆)alkynyl(C₃-C₈)cycloalkyl, -(C₂-C₆)alkynyl(C₅-C₈)cycloalkenyl, -(C₂-C₆)alkynylphenyl, -(C₂-C₆)alkynylnaphthyl, -(C₂-C₆)alkynyl(C₁₄)aryl, and -(C₂-C₆)alkynyl(C₃-C₇)heterocycle, wherein each first and second group is defined above.

Where a substituent includes a first and a second group, for example, -(C₁-C₆)alkylphenyl, wherein -(C₁-C₆)alkyl is the first group and phenyl is the second group, it is to be understood that the first group forms a bond with the Thiadiazolylpiperazine Compound's backbone and the second group is a substituent of the first group.

"Animal" includes but is not limited to, a cow, monkey, chimpanzee, baboon, horse, sheep, pig, chicken, turkey, quail, cat, dog, mouse, rat, rabbit, and guinea pig. In one embodiment, the animal is a mammal. In another embodiment the animal is a human.

The phrase "pharmaceutically acceptable salt," as used herein, is a salt formed from an acid and the basic nitrogen group of one of the Thiadiazolylpiperazine Compounds. Representative salts include, but are not limited, to sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, and pamoate (*i.e.*, 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. The term "pharmaceutically acceptable salt" also refers to a salt prepared from a Thiadiazolylpiperazine Compound having an acidic functional group, such as a carboxylic acid functional group, and a pharmaceutically acceptable inorganic or organic base. Suitable bases include, but are not limited to, hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)amine, 2-hydroxy-tert-butylamine, or tris-(hydroxymethyl)methylamine, N, N,-di-lower alkyl-N-(hydroxy lower alkyl)-amines, such as N,N,-dimethyl-N-(2-hydroxyethyl)amine, or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like.

The term "UI" means urinary incontinence.

The term "IBD" means inflammatory-bowel disease.

The term "IBS" means irritable-bowel syndrome.

The term "ALS" means amyotrophic lateral sclerosis.

The term "DMSO" means dimethyl sulfoxide.

The term "DMF" means dimethyl formamide.

The term "DCM" means dichloromethane.

The term "TEA" means triethylamine.

The term "DIEA" means di-isopropylethylamine.

The term "THF" means tetrahydrofuran.

The phrase "treatment of" and "treating" includes the amelioration or cessation of pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia,


retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression, or a symptom thereof.

The phrase "prevention of" and "preventing" includes the avoidance of the onset of pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease,
 5 parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression, or a symptom thereof.


The phrase "effective amount," when used in connection with a
 10 Thiadiazolylpiperazine Compound means an amount effective for: (a) treating or preventing pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression; or (b) inhibiting VR1,
 15 mGluR1, or mGluR5 function in a cell.

The phrase "effective amount," when used in connection with the another therapeutic agent means an amount for providing the therapeutic effect of the therapeutic agent.

When a first group is "substituted with one or more" second groups, one or
 20 more hydrogen atoms of the first group is replaced with a corresponding number of second groups. When the number of second groups is two or greater, each second group can be the same or different. In one embodiment, the number of second groups is one or two. In another embodiment, the number of second groups is one.



as used herein, means an ethylene group attached to a terminal Ar group, wherein one or each
 of two of the hydrogens of the ethylene group can optionally be substituted with an R^8 group.




as used herein, means a *n*-propylene group attached to a terminal Ar group, wherein one or each of two of the hydrogens of the propylene group can optionally be substituted with an R⁸ group.

It is to be understood that when R³ is $-(CH_2)_nOR^6$; *n* is 0; and R⁶ is -H, then R³ is unsubstituted. Similarly, it is understood that when R⁷ is $-(CH_2)_nOR^6$ or $-(CH_2)_nSR^6$; *n* is 0; and R⁶ is -H, then R⁷ is unsubstituted.

4.5 METHODS FOR MAKING THE THIADIAZOLYLPIPERAZINE COMPOUNDS

The Thiadiazolylpiperazine Compounds can be obtained using conventional organic syntheses or by the following illustrative methods shown in the scheme below:



wherein R¹, R³, R⁴, R⁵, and X are as defined above.

Compound A is reacted with Compound B in an aprotic solvent, typically with heating, to provide Compound C. Suitable aprotic solvents include, but are not limited to, DMSO, DMF, DCM, 1,2-dichloroethane, and dioxane. Compound C is isolated from the reaction mixture and purified, for example, using column chromatography or recrystallization. Compound C is then reacted with an isocyanate (R⁵-N=C=O) or isothiocyanate (R⁵-N=C=S) in a suitable solvent, at a suitable temperature, typically at room temperature, to provide the compound of Formula (I) wherein X = O or X = S, respectively. In one embodiment, the solvent is dichloromethane,


If Compound C is substituted with a hydroxyl group or thiol group, then the hydroxyl or thiol group of compound B is typically protected with a suitable protecting group before being reacted with the isocyanate or isothiocyanate. Suitable protecting groups for hydroxyl group include, but are not limited to, methyl ether, methoxymethyl ether, methoxythiomethyl ether, 2-methoxyethoxymethyl ether, bis(2-chloroethoxy)ethyl ether, tetrahydropyranyl ether, tetrahydrothiopyranyl ether, 4-methoxytetrahydropyranyl ether, methoxytetrahydrothiopyranyl ether, tetrahydrofuranyl ether, tetrahydrothiofuranyl ether, 1-ethoxyethyl ether, 1-methyl-1-methoxyethyl ether, 2-(phenylselenyl ether), *t*-butyl ether, allyl ether, benzyl ether, *o*-nitrobenzyl ether, triphenylmethyl ether, *o*-naphthylidiphenylmethyl ether, *p*-methoxydiphenylmethyl ether, 9-(9-phenyl-10-oxo)anthryl ether (tritylone), trimethylsilyl ether, isopropyl dimethylsilyl ether, *t*-butyldimethylsilyl ether, *t*-butyldiphenylsilyl ether, tribenzylsilyl ether, triisopropylsilyl ether, formate ester, acetate ester, trichloroacetate ester, phenoxyacetate ester, isobutyrate ester, pivaloate ester, adamantate ester, benzoate ester, 2,4,6-trimethyl (mesitoate) ester, methyl carbonate, 2,2,2-trichlorocarbonate, allyl carbonate, *p*-nitrophenyl carbonate, benzyl carbonate, *p*-nitrobenzyl carbonate, *S*-benzylthiocarbonate, *N*-phenylcarbamate, nitrate ester, and 2,4-dinitrophenylsulfonate ester (See, e.g., T.W. Greene, *Protective Groups in Organic Synthesis*, John Wiley-Interscience Publication, New York, (1981)).

Suitable protecting groups for a thiol group include, but are not limited to, *S*-benzyl thioether, *S-p*-methoxybenzyl thioether, *S-p*-nitrobenzyl thioether, *S*-4-picolyl thioether, *S*-2-picolyl *N*-oxide thioether, *S*-9-anthrylmethyl thioether, *S*-diphenylmethyl thioether, *S*-di(*p*-methoxyphenyl)methyl thioether, *S*-triphenylmethyl thioether, *S*-2,4-dinitrophenyl thioether, *S-t*-butyl thioether, *S*-isobutoxymethyl hemithioacetal, *S*-2-tetrahydropyranyl hemithioacetal, *S*-acetamidomethyl aminothioacetal, *S*-cyanomethyl thioether, *S*-2-nitro-1-phenylethyl thioether, *S*-2,2-bis(carboethoxy)ethyl thioether, *S*-benzoyl derivative, *S*-(*N*-ethylcarbamate), and *S*-ethyldisulfide. *Id.*

Compound A wherein R¹ is -Cl and Y is -Cl is commercially available from Aldrich Chemical Co., Milwaukee, WI, USA. Compound A wherein R¹ is -CH₃ and Y is -Cl can be prepared according to the method disclosed in U.S. Patent No. 4,145,546 to T. Brown et. al. Compound A wherein R¹ is Br and Y is Br can be made according to the procedure disclosed in DE 19712351 to R. Fuss et al. Compound A wherein R¹ is F and Y is F can be prepared by the procedure of M. Geisel et al., "Fluorination Reactions on 3, 4-dichloro-1, 2, 5-Thiadiazolylpiperazine," *Chem. Ber.* 115(6), pp. 2135-40, 1982. Compound C wherein R¹ is I can be prepared from Compound C, wherein R¹ is bromide, according to the procedure of B.S. Lee, *Bioorg. Med. Chem. Lett.*, 12:811-815 (2002).


Isocyanates R⁵-N=C=O are commercially available or preparable by reacting R⁵NH₂ with phosgene according to well-known methods (See, e.g., H. Eckert and B. Foster, *Angew. Chem. Int. Ed. Engl.*, 26:894 (1987); H. Eckert, Ger. Offen. DE 3 440 141; *Chem*

Abstr. 106, 4294d, (1987); and L. Contarca et al., *Synthesis*, 553-576 (1996)). For example an amine, R^5-NH_2 , can be reacted with triphosgene according to the scheme shown below.




Typically, a solution of triphosgene (0.3 eq) in 1,2-dichloromethane (0.3 M) is slowly added to a stirred solution of the amine (1.0 eq.) in 1, 2-dichloroethane (0.3 M) at room temperature. The reaction mixture is then stirred at room temperature for 10 min. and the temperature then raised to 70 °C. After stirring at 70 °C for 3 h., the reaction mixture is cooled to room temperature, filtered, and the filtrate concentrated to give the desired isocyanate.

Isothiocyanates $R^5-N=C=S$ are commercially available or preparable by reacting R^5NH_2 with thiophosgene as shown in the scheme below (See, e.g., *Tetrahedron Lett.*, 41(37), 7207-7209 (2000); *Synlett*, 11:1784-1786 (1999); *Heterocycles*, 32:2343-2355 (1991); *Org. Prep., Proced., Int.*, 23(6):729-734 (1991); *J. Heterocycle Chem.*, 28(4):1091-1097 (1991); *J. Fluorine Chem.*, 41(3):303-310 (1988); and *J. Med. Chem.*, 32(6):1173-1176 and 1392-1398 (1989); and *Tett. Lett.*, 42(32):5414-5416 (2001).




Alternatively, isothiocyanates $R^5-N=C=S$ can be prepared by reacting R^5NH_2 , with carbon disulfide in the presence of TEA in THF, followed by reaction with hydrogen peroxide and hydrochloric acid in water as shown in the scheme below (See, e.g., *J. Org. Chem.*, 62(13):4539-4540 (1997)).




The Thiadiazolylpiperazine Compounds can also be prepared by reacting R^5NH_2 with 4-nitrophenyl chloroformate (commercially available from Aldrich Chemical Co. of Milwaukee, WI) to provide a carbamate, Compound **D**, and then reacting Compound **D** with Compound **C** as shown in the scheme below (See, e.g., *J. Org. Chem.*, 63(23):8515-8521, (1998) and European Patent Publication No. 549 039).

The urea or thiourea nitrogen atom of R^5 of the Thiadiazolylpiperazine compounds can be alkylated using sodium hydride in an anhydrous aprotic solvent followed by reaction with an alkyl halide, such as methyl iodide.



wherein R^1 , R^3 , R^4 , and R^5 are as defined above.

The scheme below illustrates methods for making the Thiadiazolylpiperazine Compounds of formula (III) and (IV). These compounds can be prepared by reacting a Compound C with a phenethylamine, Compound E, or a phenpropylamine, Compound F, using 4-nitrophenyl chloroformate as described above.



wherein R^1 , R^2 , R^3 , R^4 , R^8 , Ar, n and p are defined above for the Thiadiazolylpiperazine Compounds of formula (III) and (IV).

In an illustrative procedure, to a 0.5M solution of 4-nitrophenyl chloroformate
5 in DCM at 0° C is added a 0.5 M solution of Compound E or Compound F (about 1 eq.) in
DCM, followed by triethylamine (about 1.1 eq.). The resulting solution is heated to about
40° C and after about 5 h the total volume of the reaction mixture is reduced by about 50
percent under reduced pressure and Compound C is added. The resulting reaction mixture is
stirred at about 40° C for about 1 h and then concentrated under reduced pressure to provide a
10 crude residue that is purified using flash chromatography (gradient elution from 100% hexanes
to 10% hexanes/ethyl acetate using a 5% gradient) to provide the corresponding compound of
formula (III) or (IV).

Certain Thiadiazolylpiperazine Compounds may have asymmetric centers and
therefore exist in different enantiomeric and diastereomeric forms. This invention relates to the
15 use of all optical isomers and stereoisomers of the Thiadiazolylpiperazine Compounds, and
mixtures thereof, and to all pharmaceutical compositions and methods of treatment that may
employ or contain them.

In addition, one or more hydrogen, carbon or other atoms of a
Thiadiazolylpiperazine Compound can be replaced by an isotope of the hydrogen, carbon or
20 other atoms. Such compounds, which are encompassed by the present invention, are useful
as research and diagnostic tools in metabolism pharmacokinetic studies and in binding
assays.

In the Thiadiazolylpiperazine Compounds the R^3 group can be on any carbon
of the Thiadiazolylpiperazine Compound's piperazine ring. In one embodiment, R^3 is a
25 group other than hydrogen and the R^3 group is attached to a carbon atom adjacent to the
nitrogen atom attached to the thiadiazolyl group. In another embodiment, R^3 is a group other
than hydrogen and the R^3 group is attached to a carbon atom adjacent to the nitrogen atom
attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally
substituted with one or two R^5 groups; or $-C(=O)-NH(CH_2)_3Ar$ group optionally substituted
30 with one or two R^8 groups.

In another embodiment, the Thiadiazolylpiperazine Compound has an R^3
group that is other than hydrogen and the carbon atom to which the R^3 group is attached has
the (R) configuration. In another embodiment, the Thiadiazolylpiperazine Compound has an
 R^3 group that is other than hydrogen and the carbon atom to which the R^3 group is attached
35 has the (S) configuration.

In another embodiment, the Thiadiazolylpiperazine Compound has an R^3
group that is other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a
nitrogen atom attached to the thiadiazolyl group, and the carbon atom to which the R^3 group

is attached has the (R) configuration. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl group, the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-(C_1-C_4)$ alkyl unsubstituted or substituted with one or more halo groups. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl, the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-CH_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl group, the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-CF_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl group, the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-CH_2CH_3$.

In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group that is other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; and the carbon atom to which the R^3 group is attached has the (R) configuration. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-(C_1-C_4)$ alkyl unsubstituted or substituted with one or more halo groups. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-CH_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-CH_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; the carbon atom to

which the R^3 group is attached has the (R) configuration, and R^3 is $-CF_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; the carbon atom to which the R^3 group is attached has the (R) configuration, and R^3 is $-CH_2CH_3$.

In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group that is other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the thiadiazolyl group, and the carbon atom to which the R^3 group is attached has the (S) configuration. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl group, the carbon atom to which the R^3 group is attached has the (S) configuration, and R^3 is $-(C_1-C_4)alkyl$ unsubstituted or substituted with one or more halo groups. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl, the carbon atom to which the R^3 group is attached has the (S) configuration, and R^3 is $-CH_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl group, the carbon atom to which the R^3 group is attached has the (S) configuration, and R^3 is $-CF_3$. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen attached to the thiadiazolyl group, the carbon atom to which the R^3 group is attached has the (S) configuration, and R^3 is $-CH_2CH_3$.

In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group that is other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; and the carbon atom to which the R^3 group is attached has the (S) configuration. In another embodiment, the Thiadiazolylpiperazine Compound has an R^3 group other than hydrogen, the R^3 group is attached to a carbon atom adjacent to a nitrogen atom attached to the $-C(=O)-NHR^5$; $-C(=S)-NHR^5$; $-C(=O)-NH(CH_2)_2Ar$ group, optionally substituted with one or two R^8 groups; or $-C(=O)-NH(CH_2)_3Ar$ group, optionally substituted with one or two R^8 groups; the carbon atom to which the R^3 group is attached has

the (S) configuration, and R³ is -(C₁-C₄)alkyl unsubstituted or substituted with one or more halo groups. In another embodiment, the Thiadiazolylpiperazine Compound has an R³ group other than hydrogen, the R³ group is attached to a carbon atom adjacent to a nitrogen atom attached to the -C(=O)-NHR⁵; -C(=S)-NHR⁵; -C(=O)-NH(CH₂)₂Ar group, optionally substituted with one or two R⁸ groups; or -C(=O)-NH(CH₂)₃Ar group, optionally substituted with one or two R⁸ groups; the carbon atom to which the R³ group is attached has the (S) configuration, and R³ is -CH₃. In another embodiment, the Thiadiazolylpiperazine Compound has an R³ group other than hydrogen, the R³ group is attached to a carbon atom adjacent to a nitrogen atom attached to the -C(=O)-NHR⁵; -C(=S)-NHR⁵; -C(=O)-NH(CH₂)₂Ar group, optionally substituted with one or two R⁸ groups; or -C(=O)-NH(CH₂)₃Ar group, optionally substituted with one or two R⁸ groups; the carbon atom to which the R³ group is attached has the (S) configuration, and R³ is -CF₃. In another embodiment, the Thiadiazolylpiperazine Compound has an R³ group other than hydrogen, the R³ group is attached to a carbon atom adjacent to a nitrogen atom attached to the -C(=O)-NHR⁵; -C(=S)-NHR⁵; -C(=O)-NH(CH₂)₂Ar group, optionally substituted with one or two R⁸ groups; or -C(=O)-NH(CH₂)₃Ar group, optionally substituted with one or two R⁸ groups; the carbon atom to which the R³ group is attached has the (S) configuration, and R³ is -CH₂CH₃.

4.6 THERAPEUTIC USES OF THE THIADIAZOLYLPIPERAZINE COMPOUNDS

In accordance with the invention, the Thiadiazolylpiperazine Compounds are administered to an animal in need of treatment or prevention of pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression.

In one embodiment, an effective amount of a Thiadiazolylpiperazine Compound can be used to treat or prevent any condition treatable or preventable by inhibiting VR1. Examples of conditions that are treatable or preventable by inhibiting VR1 include, but are not limited to, pain, UI, an ulcer, IBD, and IBS.

In another embodiment, an effective amount of a Thiadiazolylpiperazine Compound can be used to treat or prevent any condition treatable or preventable by inhibiting mGluR5. Examples of conditions that are treatable or preventable by inhibiting mGluR5

include, but are not limited to, pain, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, a pruritic condition, and psychosis.

In another embodiment, an effective amount of a Thiadiazolylpiperazine Compound can be used to treat or prevent any condition treatable or preventable by inhibiting mGluR1. Examples of conditions that are treatable or preventable by inhibiting mGluR1 include, but are not limited to, pain, UI, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, and depression.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent acute or chronic pain. Examples of pain treatable or preventable using the Thiadiazolylpiperazine Compounds include, but are not limited to, cancer pain, central pain, labor pain, myocardial infarction pain, pancreatic pain, colic pain, post-operative pain, headache pain, muscle pain, pain associated with intensive care, arthritic pain, and pain associated with a periodontal disease, including gingivitis and periodontitis.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent UI. Examples of UI treatable or preventable using the Thiadiazolylpiperazine Compounds include, but are not limited to, urge incontinence, stress incontinence, overflow incontinence, neurogenic incontinence, and total incontinence.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent an ulcer. Examples of ulcers treatable or preventable using the Thiadiazolylpiperazine Compounds include, but are not limited to, a duodenal ulcer, a gastric ulcer, a marginal ulcer, an esophageal ulcer, or a stress ulcer.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent IBD, including Crohn's disease and ulcerative colitis.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent IBS. Examples of IBS treatable or preventable using the Thiadiazolylpiperazine Compounds include, but are not limited to, spastic-colon-type IBS and constipation-predominant IBS.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent an addictive disorder, including but not limited to, an eating disorder, an impulse-control disorder, an alcohol-related disorder, a nicotine-related disorder, an amphetamine-related disorder, a cannabis-related disorder, a cocaine-related disorder, an hallucinogen-related disorder, an inhalant-related disorders, and an opioid-related disorder, all of which are further sub-classified as listed below.

Eating disorders include, but are not limited to, Bulimia Nervosa, Nonpurging Type; Bulimia Nervosa, Purging Type; Anorexia; and Eating Disorder not otherwise specified (NOS).

- 5 Impulse control disorders include, but are not limited to, Intermittent Explosive Disorder, Kleptomania, Pyromania, Pathological Gambling, Trichotillomania, and Impulse Control Disorder not otherwise specified (NOS).

- 10 Alcohol-related disorders include, but are not limited to, Alcohol-Induced Psychotic Disorder with delusions, Alcohol Abuse, Alcohol Intoxication, Alcohol Withdrawal, Alcohol Intoxication Delirium, Alcohol Withdrawal Delirium, Alcohol-Induced Persisting Dementia, Alcohol-Induced Persisting Amnesic Disorder, Alcohol Dependence, Alcohol-Induced Psychotic Disorder with hallucinations, Alcohol-Induced Mood Disorder, Alcohol-Induced Anxiety Disorder, Alcohol-Induced Sexual Dysfunction, Alcohol-Induced Sleep Disorder, Alcohol-Related Disorder not otherwise specified (NOS), Alcohol Intoxication, and Alcohol Withdrawal.

- 15 Nicotine-related disorders include, but are not limited to, Nicotine Dependence, Nicotine Withdrawal, and Nicotine-Related Disorder not otherwise specified (NOS).

- 20 Amphetamine-related disorders include, but are not limited to, Amphetamine Dependence, Amphetamine Abuse, Amphetamine Intoxication, Amphetamine Withdrawal, Amphetamine Intoxication Delirium, Amphetamine-Induced Psychotic Disorder with delusions, Amphetamine-Induced Psychotic Disorders with hallucinations, Amphetamine-Induced Mood Disorder, Amphetamine-Induced Anxiety Disorder, Amphetamine-Induced Sexual Dysfunction, Amphetamine-Induced Sleep Disorder, Amphetamine Related Disorder not otherwise specified (NOS), Amphetamine Intoxication, 25 and Amphetamine Withdrawal.

- 30 Cannabis-related disorders include, but are not limited to, Cannabis Dependence, Cannabis Abuse, Cannabis Intoxication, Cannabis Intoxication Delirium, Cannabis-Induced Psychotic Disorder with delusions, Cannabis-Induced Psychotic Disorder with hallucinations, Cannabis-Induced Anxiety Disorder, Cannabis Related Disorder not otherwise specified (NOS), and Cannabis Intoxication.

Cocaine-related disorders include, but are not limited to, Cocaine Dependence, Cocaine Abuse, Cocaine Intoxication, Cocaine Withdrawal, Cocaine Intoxication Delirium, Cocaine-Induced Psychotic Disorder with delusions, Cocaine-Induced Psychotic Disorders with hallucinations, Cocaine-Induced Mood Disorder, Cocaine-Induced Anxiety Disorder,

Cocaine-Induced Sexual Dysfunction, Cocaine-Induced Sleep Disorder, Cocaine Related Disorder not otherwise specified (NOS), Cocaine Intoxication, and Cocaine Withdrawal.

Hallucinogen-related disorders include, but are not limited to, Hallucinogen Dependence, Hallucinogen Abuse, Hallucinogen Intoxication, Hallucinogen Withdrawal, 5 Hallucinogen Intoxication Delirium, Hallucinogen-Induced Psychotic Disorder with delusions, Hallucinogen-Induced Psychotic Disorders with hallucinations, Hallucinogen-Induced Mood Disorder, Hallucinogen-Induced Anxiety Disorder, Hallucinogen-Induced Sexual Dysfunction, Hallucinogen-Induced Sleep Disorder, Hallucinogen Related Disorder not otherwise specified (NOS), Hallucinogen Intoxication, 10 and Hallucinogen Persisting Perception Disorder (Flashbacks).

Inhalant-related disorders include, but are not limited to, Inhalant Dependence, Inhalant Abuse, Inhalant Intoxication, Inhalant Intoxication Delirium, Inhalant-Induced Psychotic Disorder with delusions, Inhalant-Induced Psychotic Disorder with hallucinations, Inhalant-Induced Anxiety Disorder, Inhalant Related Disorder not otherwise specified (NOS), 15 and Inhalant Intoxication.

Opioid-related disorders include, but are not limited to, Opioid Dependence, Opioid Abuse, Opioid Intoxication, Opioid Intoxication Delirium, Opioid-Induced Psychotic Disorder with delusions, Opioid-Induced Psychotic Disorder with hallucinations, Opioid-Induced Anxiety Disorder, Opioid Related Disorder not otherwise specified (NOS), 20 Opioid Intoxication, and Opioid Withdrawal.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent Parkinson's disease and parkinsonism and the symptoms associated with Parkinson's disease and parkinsonism, including but not limited to, bradykinesia, muscular rigidity, resting tremor, and impairment of postural balance.

25 The Thiadiazolylpiperazine Compounds can be used to treat or prevent generalized anxiety or severe anxiety and the symptoms associated with anxiety, including but not limited to, restlessness; tension; tachycardia; dyspnea; depression, including chronic "neurotic" depression; panic disorder; agoraphobia and other specific phobias; eating disorders; and personality disorders.

30 The Thiadiazolylpiperazine Compounds can be used to treat or prevent epilepsy, including but not limited to, partial epilepsy, generalized epilepsy, and the symptoms associated with epilepsy, including but not limited to, simple partial seizures, jacksonian seizures, complex partial (psychomotor) seizures, convulsive seizures (grand mal or tonic-clonic seizures), petit mal (absence) seizures, and status epilepticus.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent strokes, including but not limited to, ischemic strokes and hemorrhagic strokes.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent a seizure, including but not limited to, infantile spasms, febrile seizures, and epileptic seizures.

5 The Thiadiazolylpiperazine Compounds can be used to treat or prevent a pruritic condition, including but not limited to, pruritus caused by dry skin, scabies, dermatitis, herpetiformis, atopic dermatitis, *pruritus vulvae et ani*, miliaria, insect bites, pediculosis, contact dermatitis, drug reactions, urticaria, urticarial eruptions of pregnancy, psoriasis, lichen planus, lichen simplex chronicus, exfoliative dermatitis, folliculitis, bullous
10 pemphigoid, or fiberglass dermatitis.

 The Thiadiazolylpiperazine Compounds can be used to treat or prevent psychosis, including but not limited to, schizophrenia, including paranoid schizophrenia, hebephrenic or disorganized schizophrenia, catatonic schizophrenia, undifferentiated schizophrenia, negative or deficit subtype schizophrenia, and non-deficit schizophrenia; a
15 delusional disorder, including erotomanic subtype delusional disorder, grandiose subtype delusional disorder, jealous subtype delusional disorder, persecutory subtype delusional disorder, and somatic subtype delusional disorder; and brief psychosis.

 The Thiadiazolylpiperazine Compounds can be used to treat or prevent a cognitive disorder, including but not limited to, delirium and dementia such as multi-infarct
20 dementia, dementia pugilistica, dementia caused by AIDS, and dementia caused by Alzheimer's disease.

 The Thiadiazolylpiperazine Compounds can be used to treat or prevent a memory deficiency, including but not limited to, dissociative amnesia and dissociative fugue.

 The Thiadiazolylpiperazine Compounds can be used to treat or prevent
25 restricted brain function, including but not limited to, that caused by surgery or an organ transplant, restricted blood supply to the brain, a spinal cord injury, a head injury, hypoxia, cardiac arrest, or hypoglycemia.

 The Thiadiazolylpiperazine Compounds can be used to treat or prevent Huntington's chorea.

30 The Thiadiazolylpiperazine Compounds can be used to treat or prevent ALS.

 The Thiadiazolylpiperazine Compounds can be used to treat or prevent retinopathy, including but not limited to, arteriosclerotic retinopathy, diabetic arteriosclerotic retinopathy, hypertensive retinopathy, non-proliferative retinopathy, and proliferative retinopathy.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent a muscle spasm.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent a migraine.

5 The Thiadiazolylpiperazine Compounds can be used to treat or prevent vomiting, including but not limited to, nausea vomiting, dry vomiting (retching), and regurgitation.

The Thiadiazolylpiperazine Compounds can be used to treat or prevent dyskinesia, including but not limited to, tardive dyskinesia and biliary dyskinesia.

10 The Thiadiazolylpiperazine Compounds can be used to treat or prevent depression, including but not limited to, major depression and bipolar disorder.

Applicants believe that the Thiadiazolylpiperazine Compounds are antagonists for VR1.

The invention also relates to methods for inhibiting VR1 function in a cell comprising contacting a cell capable of expressing VR1 with an effective amount of a Thiadiazolylpiperazine Compound. This method can be used *in vitro*, for example, as an assay to select cells that express VR1 and, accordingly, are useful as part of an assay to select compounds useful for treating or preventing pain, UI, an ulcer, IBD, or IBS. The method is also useful for inhibiting VR1 function in a cell *in vivo*, in an animal, a human in one embodiment, by contacting a cell, in an animal, with an effective amount of a Thiadiazolylpiperazine Compound. In one embodiment, the method is useful for treating or preventing pain in an animal. In another embodiment, the method is useful for treating or preventing UI in an animal. In another embodiment, the method is useful for treating or preventing an ulcer in an animal. In another embodiment, the method is useful for treating or preventing IBD in an animal. In another embodiment, the method is useful for treating or preventing IBS in an animal.

Examples of tissue comprising cells capable of expressing VR1 include, but are not limited to, neuronal, brain, kidney, urothelium, and bladder tissue. Methods for assaying cells that express VR1 are well known in the art.

30 Applicants believe that the Thiadiazolylpiperazine Compounds are antagonists for mGluR5.

The invention also relates to methods for inhibiting mGluR5 function in a cell comprising contacting a cell capable of expressing mGluR5 with an amount of a Thiadiazolylpiperazine Compound effective to inhibit mGluR5 function in the cell. This

method can be used *in vitro*, for example, as an assay to select cells that express mGluR5 and, accordingly, are useful as part of an assay to select compounds useful for treating or preventing pain, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, a pruritic condition, or psychosis. The method is also useful for inhibiting mGluR5 function in a cell *in vivo*, in an animal, a human in one embodiment, by contacting a cell, in an animal, with an amount of a Thiadiazolylpiperazine Compound effective to inhibit mGluR5 function in the cell. In one embodiment, the method is useful for treating or preventing pain in an animal in need thereof. In another embodiment, the method is useful for treating or preventing an addictive disorder in an animal in need thereof. In another embodiment, the method is useful for treating or preventing Parkinson's disease in an animal in need thereof. In another embodiment, the method is useful for treating or preventing parkinsonism in an animal in need thereof. In another embodiment, the method is useful for treating or preventing anxiety in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a pruritic condition in an animal in need thereof. In another embodiment, the method is useful for treating or preventing psychosis in an animal in need thereof.

Examples of cells capable of expressing mGluR5 are neuronal and glial cells of the central nervous system, particularly the brain, especially in the nucleus accumbens. Methods for assaying cells that express mGluR5 are well known in the art.

Applicants believe that the Thiadiazolylpiperazine Compounds are antagonists for mGluR1.

The invention also relates to methods for inhibiting mGluR1 function in a cell comprising contacting a cell capable of expressing mGluR1 with an amount of a Thiadiazolylpiperazine Compound effective to inhibit mGluR1 function in the cell. This method can be used *in vitro*, for example, as an assay to select cells that express mGluR1 and, accordingly, are useful as part of an assay to select compounds useful for treating or preventing pain, UI, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression. The method is also useful for inhibiting mGluR1 function in a cell *in vivo*, in an animal, a human in one embodiment, by contacting a cell, in an animal, with an amount of a Thiadiazolylpiperazine Compound effective to inhibit mGluR1 function in the cell. In one embodiment, the method is useful for treating or preventing pain in an animal in need thereof. In another embodiment, the method is useful for treating or preventing UI in an animal in need thereof. In another embodiment,

the method is useful for treating or preventing an addictive disorder in an animal in need thereof. In another embodiment, the method is useful for treating or preventing Parkinson's disease in an animal in need thereof. In another embodiment, the method is useful for treating or preventing parkinsonism in an animal in need thereof. In another embodiment, the method is useful for treating or preventing anxiety in an animal in need thereof. In another embodiment, the method is useful for treating or preventing epilepsy in an animal in need thereof. In another embodiment, the method is useful for treating or preventing stroke in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a seizure in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a pruritic condition in an animal in need thereof. In another embodiment, the method is useful for treating or preventing psychosis in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a cognitive disorder in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a memory deficit in an animal in need thereof. In another embodiment, the method is useful for treating or preventing restricted brain function in an animal in need thereof. In another embodiment, the method is useful for treating or preventing Huntington's chorea in an animal in need thereof. In another embodiment, the method is useful for treating or preventing ALS in an animal in need thereof. In another embodiment, the method is useful for treating or preventing dementia in an animal in need thereof. In another embodiment, the method is useful for treating or preventing retinopathy in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a muscle spasm in an animal in need thereof. In another embodiment, the method is useful for treating or preventing a migraine in an animal in need thereof. In another embodiment, the method is useful for treating or preventing vomiting in an animal in need thereof. In another embodiment, the method is useful for treating or preventing dyskinesia in an animal in need thereof. In another embodiment, the method is useful for treating or preventing depression in an animal in need thereof.

Examples of cells capable of expressing mGluR1 include, but are not limited to, cerebellar Purkinje neuron cells, Purkinje cell bodies (punctate), cells of spine(s) of the cerebellum; neurons and neurophil cells of olfactory-bulb glomeruli; cells of the superficial layer of the cerebral cortex; hippocampus cells; thalamus cells; superior colliculus cells; and spinal trigeminal nucleus cells. Methods for assaying cells that express mGluR1 are well known in the art.

**4.6.1 THERAPEUTIC/PROPHYLACTIC
ADMINISTRATION AND COMPOSITIONS OF THE INVENTION**

Due to their activity, the Thiadiazolylpiperazine Compounds are advantageously useful in veterinary and human medicine. As described above, the Thiadiazolylpiperazine Compounds are useful for treating or preventing pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression in an animal in need thereof.

10 When administered to an animal, the Thiadiazolylpiperazine Compounds are administered as a component of a composition that comprises a pharmaceutically acceptable carrier or excipient. The present compositions, which comprise a Thiadiazolylpiperazine Compound, can be administered orally. The Thiadiazolylpiperazine Compounds of the invention can also be administered by any other convenient route, for example, by infusion or
15 bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral, rectal, and intestinal mucosa, *etc.*) and can be administered together with another biologically active agent. Administration can be systemic or local. Various delivery systems are known, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, capsules, *etc.*, and can be used to administer the Thiadiazolylpiperazine Compound.

20 Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, by inhalation, or topical, particularly to the ears, nose, eyes, or skin. The mode of administration is left to the discretion of the practitioner. In most instances, administration will result in the release of
25 the Thiadiazolylpiperazine Compounds into the bloodstream.

 In specific embodiments, it can be desirable to administer the Thiadiazolylpiperazine Compounds locally. This can be achieved, for example, and not by way of limitation, by local infusion during surgery, topical application, *e.g.*, in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a
30 suppository or enema, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.

 In certain embodiments, it can be desirable to introduce the Thiadiazolylpiperazine Compounds into the central nervous system or gastrointestinal tract by any suitable route, including intraventricular, intrathecal, and epidural injection, and

enema. Intraventricular injection can be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir.

Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent, or via perfusion in a fluorocarbon or synthetic pulmonary surfactant. In certain embodiments, the Thiadiazolylpiperazine Compounds can be formulated as a suppository, with traditional binders and excipients such as triglycerides.

In another embodiment, the Thiadiazolylpiperazine Compounds can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990) and Treat *et al.*, *Liposomes in the Therapy of Infectious Disease and Cancer* 317-327 and 353-365 (1989)).

In yet another embodiment, the Thiadiazolylpiperazine Compounds can be delivered in a controlled-release system or sustained-release system (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)). Other controlled- or sustained-release systems discussed in the review by Langer, *Science* 249:1527-1533 (1990) can be used. In one embodiment, a pump can be used (Langer, *Science* 249:1527-1533 (1990); Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald *et al.*, *Surgery* 88:507 (1980); and Saudek *et al.*, *N. Engl. J. Med.* 321:574 (1989)).

In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release* (Langer and Wise eds., 1974); *Controlled Drug Bioavailability, Drug Product Design and Performance* (Smolen and Ball eds., 1984); Ranger and Peppas, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); Levy *et al.*, *Science* 228:190 (1985); During *et al.*, *Ann. Neurol.* 25:351 (1989); and Howard *et al.*, *J. Neurosurg.* 71:105 (1989)).

In yet another embodiment, a controlled- or sustained-release system can be placed in proximity of a target of the Thiadiazolylpiperazine Compounds, e.g., the spinal column, brain, or gastrointestinal tract, thus requiring only a fraction of the systemic dose.

The present compositions can optionally comprise a suitable amount of a pharmaceutically acceptable excipient so as to provide the form for proper administration to the animal.

Such pharmaceutical excipients can be liquids, such as water and oils, including those of petroleum, animal, vegetable, or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. The pharmaceutical excipients can be saline, gum acacia, gelatin, starch paste, talc, keratin, colloidal silica, urea and the like. In addition, auxiliary, stabilizing, thickening, lubricating, and coloring agents can be used. In

one embodiment, the pharmaceutically acceptable excipients are sterile when administered to an animal. Water is a particularly useful excipient when the Thiadiazolylpiperazine Compound is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid excipients, particularly for injectable solutions. Suitable pharmaceutical excipients also include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The present compositions, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

10 The present compositions can take the form of solutions, suspensions, emulsion, tablets, pills, pellets, capsules, capsules containing liquids, powders, sustained-release formulations, suppositories, emulsions, aerosols, sprays, suspensions, or any other form suitable for use. In one embodiment, the composition is in the form of a capsule (see *e.g.*, U.S. Patent No. 5,698,155). Other examples of suitable pharmaceutical excipients are described in *Remington's Pharmaceutical Sciences* 1447-1676 (Alfonso R. Gennaro ed., 19th ed. 1995), incorporated herein by reference.

 In one embodiment, the Thiadiazolylpiperazine Compounds are formulated in accordance with routine procedures as a composition adapted for oral administration to human beings. Compositions for oral delivery can be in the form of tablets, lozenges, aqueous or oily suspensions, granules, powders, emulsions, capsules, syrups, or elixirs, for example. Orally administered compositions can contain one or more agents, for example, sweetening agents such as fructose, aspartame or saccharin; flavoring agents such as peppermint, oil of wintergreen, or cherry; coloring agents; and preserving agents, to provide a pharmaceutically palatable preparation. Moreover, where in tablet or pill form, the compositions can be coated to delay disintegration and absorption in the gastrointestinal tract thereby providing a sustained action over an extended period of time. Selectively permeable membranes surrounding an osmotically active driving compound are also suitable for orally administered compositions. In these latter platforms, fluid from the environment surrounding the capsule is imbibed by the driving compound, which swells to displace the agent or agent composition through an aperture. These delivery platforms can provide an essentially zero order delivery profile as opposed to the spiked profiles of immediate release formulations. A time-delay material such as glycerol monostearate or glycerol stearate can also be used. Oral compositions can include standard excipients such as mannitol, lactose, starch, magnesium

stearate, sodium saccharin, cellulose, and magnesium carbonate. In one embodiment, the excipients are of pharmaceutical grade.

In another embodiment, the Thiadiazolylpiperazine Compounds can be formulated for intravenous administration. Typically, compositions for intravenous administration comprise sterile isotonic aqueous buffer. Where necessary, the compositions can also include a solubilizing agent. Compositions for intravenous administration can optionally include a local anesthetic such as lidocaine to lessen pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampule or sachette indicating the quantity of active agent. Where the Thiadiazolylpiperazine Compounds are to be administered by infusion, they can be dispensed, for example, with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the Thiadiazolylpiperazine Compounds are administered by injection, an ampule of sterile water for injection or saline can be provided so that the ingredients can be mixed prior to administration.

The Thiadiazolylpiperazine Compounds can be administered by controlled-release or sustained-release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; 4,008,719; 5,674,533; 5,059,595; 5,591,767; 5,120,548; 5,073,543; 5,639,476; 5,354,556; and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide controlled- or sustained-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled- or sustained-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled- or sustained-release.

Controlled- or sustained-release pharmaceutical compositions can have a common goal of improving drug therapy over that achieved by their non-controlled or non-sustained counterparts. In one embodiment, a controlled- or sustained-release composition comprises a minimal amount of a Thiadiazolylpiperazine Compound to cure or control the condition in a minimum amount of time. Advantages of controlled- or sustained-release

compositions include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled- or sustained-release compositions can favorably affect the time of onset of action or other characteristics, such as blood levels of the Thiadiazolylpiperazine Compound, and can thus reduce the occurrence of adverse side effects.

Controlled- or sustained-release compositions can initially release an amount of a Thiadiazolylpiperazine Compound that promptly produces the desired therapeutic or prophylactic effect, and gradually and continually release other amounts of the Thiadiazolylpiperazine Compound to maintain this level of therapeutic or prophylactic effect over an extended period of time. To maintain a constant level of the Thiadiazolylpiperazine Compound in the body, the Thiadiazolylpiperazine Compound can be released from the dosage form at a rate that will replace the amount of Thiadiazolylpiperazine Compound being metabolized and excreted from the body. Controlled- or sustained-release of an active ingredient can be stimulated by various conditions, including but not limited to, changes in pH, changes in temperature, concentration or availability of enzymes, concentration or availability of water, or other physiological conditions or compounds.

The amount of the Thiadiazolylpiperazine Compound that is effective in the treatment or prevention of pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression and can be determined by standardinical techniques. In addition, *in vitro* or *in vivo* assays can optionally be employed to help identify optimal dosage ranges. The precise dose to be employed will also depend on the route of administration, and the seriousness of the condition being treated and should be decided according to the judgment of the practitioner and each patient's circumstances in view of, e.g., published clinical studies. Suitable effective dosage amounts, however, range from about 10 micrograms to about 2500 milligrams about every 4 h, although they are typically about 100 mg or less. In one embodiment, the effective dosage amount ranges from about 0.01 milligrams to about 100 milligrams of a Thiadiazolylpiperazine Compound about every 4 h, in another embodiment, about 0.020 milligrams to about 50 milligrams about every 4 h, and in another embodiment, about 0.025 milligrams to about 20 milligrams about every 4 h. The effective dosage amounts described herein refer to total amounts administered; that is, if more than one

Thiadiazolylpiperazine Compound is administered, the effective dosage amounts correspond to the total amount administered.

Where a cell capable of expressing VR1, mGluR5, or mGluR1 is contacted with a Thiadiazolylpiperazine Compound *in vitro*, the amount effective for inhibiting the receptor function in a cell will typically range from about 0.01 $\mu\text{g/L}$ to about 5 mg/L, in one embodiment, from about 0.01 $\mu\text{g/L}$ to about 2.5 mg/L, in another embodiment, from about 0.01 $\mu\text{g/L}$ to about 0.5 mg/L, and in another embodiment, from about 0.01 $\mu\text{g/L}$ to about 0.25 mg/L of a solution or suspension of a pharmaceutically acceptable carrier or excipient. In one embodiment, the volume of solution or suspension is from about 1 μL to about 1 mL. In another embodiment, the volume of solution or suspension is about 200 μL .

Where a cell capable of expressing VR1, mGluR5, or mGluR1 is contacted with a Thiadiazolylpiperazine Compound *in vivo*, the amount effective for inhibiting the receptor function in a cell will typically range from about 0.01 mg to about 100 mg/kg of body weight per day, in one embodiment, from about 0.1 mg to about 50 mg/kg body weight per day, and in another embodiment, from about 1 mg to about 20 mg/kg of body weight per day.

The Thiadiazolylpiperazine Compounds can be assayed *in vitro* or *in vivo* for the desired therapeutic or prophylactic activity prior to use in humans. Animal model systems can be used to demonstrate safety and efficacy.

The present methods for treating or preventing pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression in an animal in need thereof can further comprise administering to the animal being administered a Thiadiazolylpiperazine Compound another therapeutic agent. In one embodiment, the other therapeutic agent is administered in an effective amount.

The present methods for inhibiting VR1 function in a cell capable of expressing VR1 can further comprise contacting the cell with an effective amount of another therapeutic agent.

The present methods for inhibiting mGluR5 function in a cell capable of expressing mGluR5 can further comprise contacting the cell with an effective amount of another therapeutic agent.

The present methods for inhibiting mGluR1 function in a cell capable of expressing mGluR1 can further comprise contacting the cell with an effective amount of another therapeutic agent.

The other therapeutic agent includes, but is not limited to, an opioid agonist, a
5 non-opioid analgesic, a non-steroidal anti-inflammatory agent, an antimigraine agent, a Cox-II inhibitor, an antiemetic, a β -adrenergic blocker, an anticonvulsant, an antidepressant, a Ca²⁺-channel blocker, an anticancer agent, an agent for treating or preventing UI, an agent for treating or preventing an ulcer, an agent for treating or preventing IBD, an agent for treating or preventing IBS, an agent for treating addictive disorder, an agent for treating
10 Parkinson's disease and parkinsonism, an agent for treating anxiety, an agent for treating epilepsy, an agent for treating a stroke, an agent for treating a seizure, an agent for treating a pruritic condition, an agent for treating psychosis, an agent for treating Huntington's chorea, an agent for treating ALS, an agent for treating a cognitive disorder, an agent for treating a migraine, an agent for treating vomiting, an agent for treating dyskinesia, or an agent for
15 treating depression, and mixtures thereof.

Effective amounts of the other therapeutic agents are well known to those skilled in the art. However, it is well within the skilled artisan's purview to determine the other therapeutic agent's optimal effective-amount range. In one embodiment of the invention, where another therapeutic agent is administered to an animal, the effective amount
20 of the Thiadiazolylpiperazine Compound is less than its effective amount would be where the other therapeutic agent is not administered. In this case, without being bound by theory, it is believed that the Thiadiazolylpiperazine Compounds and the other therapeutic agent act synergistically to treat or prevent pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition,
25 psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression.

Examples of useful opioid agonists include, but are not limited to, alfentanil, allylprodine, alphaprodine, anileridine, benzylmorphine, bezitramide, buprenorphine,
30 butorphanol, clonitazene, codeine, desomorphine, dextromoramide, dezocine, diampromide, diamorphine, dihydrocodeine, dihydromorphine, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, eptazocine, ethoheptazine, ethylmethylthiambutene, ethylmorphine, etonitazene fentanyl, heroin, hydrocodone, hydromorphone, hydroxypethidine, isomethadone, ketobemidone, levorphanol,

levophenacymorphan, lofentanil, meperidine, meptazinol, metazocine, methadone, metopon, morphine, myrophine, nalbuphine, narceine, nicomorphine, norlevorphanol, normethadone, nalorphine, normorphine, norpipanone, opium, oxycodone, oxymorphone, papaveretum, pentazocine, phenadoxone, phenomorphan, phenazocine, phenoperidine, piminodine, piritramide, proheptazine, promedol, properidine, propiram, propoxyphene, sufentanil, tilidine, tramadol, pharmaceutically acceptable salts thereof, and mixtures thereof.

In certain embodiments, the opioid agonist is selected from codeine, hydromorphone, hydrocodone, oxycodone, dihydrocodeine, dihydromorphine, morphine, tramadol, oxymorphone, pharmaceutically acceptable salts thereof, and mixtures thereof.

Examples of useful non-opioid analgesics include non-steroidal anti-inflammatory agents, such as aspirin, ibuprofen, diclofenac, naproxen, benoxaprofen, flurbiprofen, fenoprofen, flubufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pramoprofen, muprofen, trioxaprofen, suprofen, aminoprofen, tiaprofenic acid, fluprofen, bucloxic acid, indomethacin, sulindac, tolmetin, zomepirac, tiopinac, zidometacin, acetaminophen, fentiazac, clidanac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid, tolfenamic acid, diflunisal, flufenisal, piroxicam, sudoxicam, isoxicam, and pharmaceutically acceptable salts thereof, and mixtures thereof. Other suitable non-opioid analgesics include the following, non-limiting, chemical classes of analgesic, antipyretic, nonsteroidal anti-inflammatory drugs: salicylic acid derivatives, including aspirin, sodium salicylate, choline magnesium trisalicylate, salsalate, diflunisal, salicylsalicylic acid, sulfasalazine, and olsalazine; para-aminophenol derivatives including acetaminophen and phenacetin; indole and indene acetic acids, including indomethacin, sulindac, and etodolac; heteroaryl acetic acids, including tolmetin, diclofenac, and ketorolac; anthranilic acids (fenamates), including mefenamic acid and meclofenamic acid; enolic acids, including oxicams (piroxicam, tenoxicam), and pyrazolidinediones (phenylbutazone, oxyphenbutazone); and alkanones, including nabumetone. For a more detailed description of the NSAIDs, see Paul A. Insel, *Analgesic-Antipyretic and Anti-inflammatory Agents and Drugs Employed in the Treatment of Gout*, in Goodman & Gilman's *The Pharmacological Basis of Therapeutics* 617-57 (Perry B. Molinoff and Raymond W. Ruddon eds., 9th ed 1996) and Glen R. Hanson, *Analgesic, Antipyretic and Anti-Inflammatory Drugs in Remington: The Science and Practice of Pharmacy Vol II* 1196-1221 (A.R. Gennaro ed. 19th ed. 1995) which are hereby incorporated by reference in their entireties.

Examples of useful Cox-II inhibitors and 5-lipoxygenase inhibitors, as well as combinations thereof, are described in U.S. Patent No. 6,136,839, which is hereby

incorporated by reference in its entirety. Examples of useful Cox-II inhibitors include, but are not limited to, rofecoxib and celecoxib.

Examples of useful antimigraine agents include, but are not limited to, alpiropride, dihydroergotamine, dolasetron, ergocornine, ergocorninine, ergocryptine, ergot,
5 ergotamine, flumetrigone acetate, fonazine, lisuride, lomerizine, methysergide oxetorone, pizotyline, and mixtures thereof.

The other therapeutic agent can also be an agent useful for reducing any potential side effects of a Thiadiazolylpiperazine Compounds. For example, the other therapeutic agent can be an antiemetic agent. Examples of useful antiemetic agents include,
10 but are not limited to, metoclopramide, domperidone, prochlorperazine, promethazine, chlorpromazine, trimethobenzamide, ondansetron, granisetron, hydroxyzine, acetylcholine monoethanolamine, alizapride, azasetron, benzquinamide, biantanautine, bromopride, buclizine, clemopride, cyclizine, dimenhydrinate, diphenidol, dolasetron, meclizine, methallatal, metopimazine, nabilone, oxypendyl, pipamazine, scopolamine, sulpiride,
15 tetrahydrocannabinol, thiethylperazine, thioproperazine, tropisetron, and mixtures thereof.

Examples of useful β -adrenergic blockers include, but are not limited to, acebutolol, alprenolol, amosulablol, arotinolol, atenolol, befunolol, betaxolol, bevantolol, bisoprolol, bopindolol, bucumolol, bufetolol, bufuralol, bunitrolol, bupranolol, butidine hydrochloride, butofilolol, carazolol, carteolol, carvedilol, celiprolol, cetamolol, cloranolol,
20 dilevalol, epanolol, esmolol, indenolol, labetalol, levobunolol, mepindolol, metipranolol, metoprolol, moprolool, nadolol, nadoxolol, nebivolol, nifenalol, nipradilol, oxprenolol, penbutolol, pindolol, practolol, pronethalol, propranolol, sotalol, sulfinalol, talinolol, tertatolol, tilisolol, timolol, toliprolol, and xibenolol.

Examples of useful anticonvulsants include, but are not limited to,
25 acetylpheneturide, albutoin, aloxidone, aminoglutethimide, 4-amino-3-hydroxybutyric acid, atrolactamide, beclamide, buramate, calcium bromide, carbamazepine, cinromide, clomethiazole, clonazepam, decimemide, diethadione, dimethadione, doxinitroin, eterobarb, ethadione, ethosuximide, ethotoin, felbamate, fluoresone, gabapentin, 5-hydroxytryptophan, lamotrigine, magnesium bromide, magnesium sulfate, mephentoin, mephobarbital,
30 metharbital, methetoin, methsuximide, 5-methyl-5-(3-phenanthryl)-hydantoin, 3-methyl-5-phenylhydantoin, narcobarbital, nimetazepam, nitrazepam, oxcarbazepine, paramethadione, phenacetamide, phenetharbital, pheneturide, phenobarbital, phenisuximide, phenylmethylbarbituric acid, phenytoin, phethenylate sodium, potassium bromide, pregabalin, primidone, progabide, sodium bromide, solanum, strontium bromide,

suclofenide, sulthiame, tetrantoin, tiagabine, topiramate, trimethadione, valproic acid, valpromide, vigabatrin, and zonisamide.

Examples of useful antidepressants include, but are not limited to, binedaline, caroxazone, citalopram, dimethazan, fencamine, indalpine, indeloxazine hydrochloride, nefopam, nomifensine, oxitriptan, oxypertine, paroxetine, sertraline, thiazesim, trazodone, benmoxine, iproclozide, iproniazid, isocarboxazid, nialamide, octamoxin, phenelzine, cotinine, rolicyprine, rolipram, maprotiline, metralindole, mianserin, mirtazepine, adinazolam, amitriptyline, amitriptylinoxide, amoxapine, butriptyline, clomipramine, demexiptiline, desipramine, dibenzepin, dimetacrine, dothiepin, doxepin, fluacizine, imipramine, imipramine N-oxide, iprindole, lofepramine, melitracen, metapramine, nortriptyline, noxiptilin, opipramol, pizotiline, propizepine, protriptyline, quinupramine, tianeptine, trimipramine, adrafinil, benactyzine, bupropion, butacetin, dioxadrol, duloxetine, etoperidone, febarbamate, femoxetine, fentpentadiol, fluoxetine, fluvoxamine, hematoporphyrin, hypericin, levophacetoperane, medifoxamine, milnacipran, minaprine, moclobemide, nefazodone, oxaflozane, piberaline, prolintane, pyrisuccideanol, ritanserin, roxindole, rubidium chloride, sulpiride, tandospirone, thozalinone, tofenacin, toloxatone, tranlycypromine, L-tryptophan, venlafaxine, viloxazine, and zimelidine.

Examples of useful Ca²⁺-channel blockers include, but are not limited to, bepridil, clentiazem, diltiazem, fendiline, gallopamil, mibefradil, prenylamine, semotiadil, terodiline, verapamil, amlodipine, aranidipine, barnidipine, benidipine, cilnidipine, efonidipine, elgodipine, felodipine, isradipine, lacidipine, lercanidipine, manidipine, nicardipine, nifedipine, nilvadipine, nimodipine, nisoldipine, nitrendipine, cinnarizine, flunarizine, lidoflazine, lomerizine, bencyclane, etafenone, fantofarone, and perhexiline.

Examples of useful anticancer agents include, but are not limited to, acivicin, aclarubicin, acodazole hydrochloride, acronine, adozelesin, aldesleukin, altretamine, ambomycin, ametantrone acetate, aminogluthethimide, amsacrine, anastrozole, anthramycin, asparaginase, asperlin, azacitidine, azetepa, azotomycin, batimastat, benzodepa, bicalutamide, bisantrene hydrochloride, bisnafide dimesylate, bizelesin, bleomycin sulfate, brequinar sodium, broprimine, busulfan, cactinomycin, calusterone, caracemide, carbetimer, carboplatin, carmustine, carubicin hydrochloride, carzelesin, cedefingol, chlorambucil, cirolemycin, cisplatin, cladribine, crisnatol mesylate, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunorubicin hydrochloride, decitabine, dexormaplatin, dezaguanine, dezaguanine mesylate, diaziquone, docetaxel, doxorubicin, doxorubicin hydrochloride, droloxifene, droloxifene citrate, dromostanolone propionate, duazomycin,

edatrexate, eflornithine hydrochloride, elsamitrucin, enloplatin, enpromate, epipropidine, epirubicin hydrochloride, erbulozole, esorubicin hydrochloride, estramustine, estramustine phosphate sodium, etanidazole, etoposide, etoposide phosphate, etoprine, fadrozole hydrochloride, fazarabine, fenretinide, floxuridine, fludarabine phosphate, fluorouracil, 5 flurocitabine, fosquidone, fostriecin sodium, gemcitabine, gemcitabine hydrochloride, hydroxyurea, idarubicin hydrochloride, ifosfamide, ilmofofosine, interleukin II (including recombinant interleukin II or rIL2), interferon alfa-2a, interferon alfa-2b, interferon alfa-n1 , interferon alfa-n3, interferon beta-I a, interferon gamma-I b, iproplatin, irinotecan hydrochloride, lanreotide acetate, letrozole, leuprolide acetate, liarozole hydrochloride, 10 lometrexol sodium, lomustine, losoxantrone hydrochloride, masoprocol, maytansine, mechlorethamine hydrochloride, megestrol acetate, melengestrol acetate, melphalan, menogaril, mercaptopurine, methotrexate, methotrexate sodium, metoprine, meturedopa, mitindomide, mitocarcin, mitocromin, mitogillin, mitomalcin, mitomycin, mitosper, mitotane, mitoxantrone hydrochloride, mycophenolic acid, nocodazole, nogalamycin, 15 ormaplatin, oxisuran, paclitaxel, pegaspargase, peliomycin, pentamustine, peplomycin sulfate, perfosfamide, pipobroman, pipo sulfan, piroxantrone hydrochloride, plicamycin, plomestane, porfimer sodium, porfiromycin, prednimustine, procarbazine hydrochloride, puromycin, puromycin hydrochloride, pyrazofurin, riboprine, rogletimide, safingol, safingol hydrochloride, semustine, simtrazene, sparfosate sodium, sparsomycin, spirogermanium 20 hydrochloride, spiromustine, spiroplatin, streptonigrin, streptozotocin, sulofenur, talisomycin, tecogalan sodium, tegafur, teloxantrone hydrochloride, temoporphin, teniposide, teroxirone, testolactone, thiamiprine, thioguanine, thiotepa, tiazofurin, tirapazamine, toremifene citrate, trestolone acetate, tricinbine phosphate, trimetrexate, trimetrexate glucuronate, triptorelin, tubulozole hydrochloride, uracil mustard, uredepa, vapreotide, verteporfin, vinblastine 25 sulfate, vincristine sulfate, vindesine, vindesine sulfate, vinepidine sulfate, vinglycinate sulfate, vinleurosine sulfate, vinorelbine tartrate, vinrosidine sulfate, vinzolidine sulfate, vorozole, zeniplatin, zinostatin, zorubicin hydrochloride.

Examples of other anti-cancer drugs include, but are not limited to, 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; 30 adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators;

- apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine;
 atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin;
 azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists;
 benzochlorins; benzoylstauosporine; beta lactam derivatives; beta-alethine; betaclamycin B;
 5 betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide;
 bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine;
 calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine;
 carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage
 derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B;
 10 cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine;
 clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4;
 combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8;
 cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplata; cypemycin;
 cytarabine ocfosfate; cytolytic factor; cytostatin; daclicimab; decitabine; dehydrodidemnin B;
 15 deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziqune;
 didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-;
 dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine;
 droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab;
 eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen
 20 agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole;
 fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone;
 fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin;
 fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase
 inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene
 25 bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofofosine;
 ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth
 factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane;
 iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B;
 itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin;
 30 lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte
 alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear
 polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds;
 lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin;
 loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine;

- mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin
- 5 fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip;
- 10 naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; odansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives;
- 15 palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds;
- 20 platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl
- 25 protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain
- 30 antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans;

tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; tricyribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer.

Examples of useful therapeutic agents for treating or preventing UI include, but are not limited to, propantheline, imipramine, hyoscyamine, oxybutynin, and dicyclomine.

Examples of useful therapeutic agents for treating or preventing an ulcer include, antacids such as aluminum hydroxide, magnesium hydroxide, sodium bicarbonate, and calcium bicarbonate; sucralfate; bismuth compounds such as bismuth subsalicylate and bismuth subcitrate; H₂ antagonists such as cimetidine, ranitidine, famotidine, and nizatidine; H⁺, K⁺ - ATPase inhibitors such as omeprazole, lansoprazole, and lansoprazole; carbenoxolone; misoprostol; and antibiotics such as tetracycline, metronidazole, timidazole, clarithromycin, and amoxicillin.

Examples of useful therapeutic agents for treating or preventing IBD include, but are not limited to, anticholinergic drugs; diphenoxylate; loperamide; deodorized opium tincture; codeine; broad-spectrum antibiotics such as metronidazole; sulfasalazine; olsalazine; mesalamine; prednisone; azathioprine; mercaptopurine; and methotrexate.

Examples of useful therapeutic agents for treating or preventing IBS include, but are not limited to, propantheline; muscarine receptor antagonists such as pirenzapine, methoctramine, ipratropium, tiotropium, scopolamine, methscopolamine, homatropine, homatropine methylbromide, and methantheline; and antidiarrheal drugs such as diphenoxylate and loperamide.

Examples of useful therapeutic agents for treating or preventing an addictive disorder include, but are not limited to, methadone, desipramine, amantadine, fluoxetine, buprenorphine, an opiate agonist, 3-phenoxypyridine, levomethadyl acetate hydrochloride, and serotonin antagonists.

Examples of useful therapeutic agents for treating or preventing Parkinson's disease and parkinsonism include, but are not limited to, carbidopa/levodopa, pergolide, bromocriptine, ropinirole, pramipexole, entacapone, tolcapone, selegiline, amantadine, and trihexyphenidyl hydrochloride.

- 5 Examples of useful therapeutic agents for treating or preventing anxiety include, but are not limited to, benzodiazepines, such as alprazolam, brotizolam, chlordiazepoxide, clobazam, clonazepam, clorazepate, demoxepam, diazepam, estazolam, flumazenil, flurazepam, halazepam, lorazepam, midazolam, nitrazepam, nordazepam, oxazepam, prazepam, quazepam, temazepam, and triazolam; non-benzodiazepine agents, 10 such as buspirone, gepirone, ipsapirone, tiospirone, zolpicone, zolpidem, and zaleplon; tranquilizers, such as barbituates, *e.g.*, amobarbital, aprobarbital, butabarbital, butalbital, mephobarbital, methohexital, pentobarbital, phenobarbital, secobarbital, and thiopental; and propanediol carbamates, such as meprobamate and tybamate.

- Examples of useful therapeutic agents for treating or preventing epilepsy 15 include, but are not limited to, carbamazepine, ethosuximide, gabapentin, lamotrigine, phenobarbital, phenytoin, primidone, valproic acid, trimethadione, benzodiazepines, gabapentin, lamotrigine, γ -vinyl GABA, acetazolamide, and felbamate.

- Examples of useful therapeutic agents for treating or preventing stroke include, but are not limited to, anticoagulants such as heparin, agents that break up clots such 20 as streptokinase or tissue plasminogen activator, agents that reduce swelling such as mannitol or corticosteroids, and acetylsalicylic acid.

- Examples of useful therapeutic agents for treating or preventing a seizure include, but are not limited to, carbamazepine, ethosuximide, gabapentin, lamotrigine, phenobarbital, phenytoin, primidone, valproic acid, trimethadione, benzodiazepines, 25 gabapentin, lamotrigine, γ -vinyl GABA, acetazolamide, and felbamate.

- Examples of useful therapeutic agents for treating or preventing a pruritic condition include, but are not limited to, naltrexone; nalmeferene; danazol; tricyclics such as amitriptyline, imipramine, and doxepin; antidepressants such as those given below, menthol; camphor; phenol; pramoxine; capsaicin; tar; steroids; and antihistamines.

- 30 Examples of useful therapeutic agents for treating or preventing psychosis include, but are not limited to, phenothiazines such as chlorpromazine hydrochloride, mesoridazine besylate, and thioridazine hydrochloride; thioxanthenes such as chlorprothixene and thiothixene hydrochloride; clozapine; risperidone; olanzapine;

quetiapine; quetiapine fumarate; haloperidol; haloperidol decanoate; loxapine succinate; molindone hydrochloride; pimozide; and ziprasidone.

Examples of useful therapeutic agents for treating or preventing Huntington's chorea include, but are not limited to, haloperidol and pimozide.

5 Examples of useful therapeutic agents for treating or preventing ALS include, but are not limited to, baclofen, neurotrophic factors, riluzole, tizanidine, benzodiazepines such as clonazepam and dantrolene.

10 Examples of useful therapeutic agents for treating or preventing cognitive disorders include, but are not limited to, agents for treating or preventing dementia such as tacrine; donepezil; ibuprofen; antipsychotic drugs such as thioridazine and haloperidol; and antidepressant drugs such as those given below.

 Examples of useful therapeutic agents for treating or preventing a migraine include, but are not limited to, sumatriptan; methysergide; ergotamine; caffeine; and beta-blockers such as propranolol, verapamil, and divalproex.

15 Examples of useful therapeutic agents for treating or preventing vomiting include, but are not limited to, 5-HT₃ receptor antagonists such as ondansetron, dolasetron, granisetron, and tropisetron; dopamine receptor antagonists such as prochlorperazine, thiethylperazine, chlorpromazine, metoclopramide, and domperidone; glucocorticoids such as dexamethasone; and benzodiazepines such as lorazepam and alprazolam.

20 Examples of useful therapeutic agents for treating or preventing dyskinesia include, but are not limited to, reserpine and tetrabenazine.

 Examples of useful therapeutic agents for treating or preventing depression include, but are not limited to, tricyclic antidepressants such as amitriptyline, amoxapine, bupropion, clomipramine, desipramine, doxepin, imipramine, maprotiline, nefazadone, 25 nortriptyline, protriptyline, trazodone, trimipramine, and venlafaxine; selective serotonin reuptake inhibitors such as fluoxetine, fluvoxamine, paroxetine, and setraline; monoamine oxidase inhibitors such as isocarboxazid, pargyline, phenelzine, and tranylcypromine; and psychostimulants such as dextroamphetamine and methylphenidate.

 A Thiadiazolylpiperazine Compound and the other therapeutic agent can act 30 additively or, in one embodiment, synergistically. In one embodiment, a Thiadiazolylpiperazine Compound is administered concurrently with another therapeutic agent. In one embodiment, a composition comprising an effective amount of a Thiadiazolylpiperazine Compound and an effective amount of another therapeutic agent can be administered. Alternatively, a composition comprising an effective amount of a

Thiadiazolylpiperazine Compound and a different composition comprising an effective amount of another therapeutic agent can be concurrently administered. In another embodiment, an effective amount of a Thiadiazolylpiperazine Compound is administered prior or subsequent to administration of an effective amount of another therapeutic agent. In this embodiment, the Thiadiazolylpiperazine Compound is administered while the other therapeutic agent exerts its therapeutic effect, or the other therapeutic agent is administered while the Thiadiazolylpiperazine Compound exerts its preventative or therapeutic effect for treating or preventing pain, UI [pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression.

4.6.2 KITS

The invention encompasses kits that can simplify the administration of a Thiadiazolylpiperazine Compound to an animal.

A typical kit of the invention comprises a unit dosage form of a Thiadiazolylpiperazine Compound. In one embodiment, the unit dosage form is a container, which can be sterile, containing an effective amount of a Thiadiazolylpiperazine Compound and a pharmaceutically acceptable carrier or excipient. The kit can further comprise a label or printed instructions instructing the use of the Thiadiazolylpiperazine Compound to treat pain, UI, an ulcer, IBD, IBS, an addictive disorder, Parkinson's disease, parkinsonism, anxiety, epilepsy, stroke, a seizure, a pruritic condition, psychosis, a cognitive disorder, a memory deficit, restricted brain function, Huntington's chorea, ALS, dementia, retinopathy, a muscle spasm, a migraine, vomiting, dyskinesia, or depression. The kit can also further comprise a unit dosage form of another therapeutic agent, for example, a container containing an effective amount of the other therapeutic agent. In one embodiment, the kit comprises a container containing an effective amount of a Thiadiazolylpiperazine Compound and an effective amount of another therapeutic agent. Examples of other therapeutic agents include, but are not limited to, those listed above.

Kits of the invention can further comprise a device that is useful for administering the unit dosage forms. Examples of such a device includes, but are not limited to, a syringe, a drip bag, a patch, an inhaler, and an enema bag.

The following examples are set forth to assist in understanding the invention and should not, of course, be construed as specifically limiting the invention described and


claimed herein. Such variations of the invention, including the substitution of all equivalents now known or later developed, which would be within the purview of those skilled in the art, and changes in formulation or minor changes in experimental design, are to be considered to fall within the scope of the invention incorporated herein.


5. EXAMPLES

Example 1 relates to the synthesis of illustrative Thiadiazolypiperazine Compounds.


Example 1: Synthesis of compounds AAV, AAW, and ABA.

Scheme A






ABA



AAV



AAW

To compound **1b** (1.2 g, 6.45 mmol) dissolved in 4 mL of DMSO was added DIEA (1.12 mL, 6.45 mmol), followed by 3, 4-dichloro-1, 2, 5-thiadiazolylpiperazine **1a** (500 mg, 3.23 mmol) and the resulting reaction mixture allowed to stir at 100 °C for 16 hrs.

The reaction mixture was then cooled to room temperature and water and ethyl acetate were added to the reaction mixture. The ethyl acetate phase and an aqueous phase were then separated and the aqueous phase was extracted twice with ethyl acetate; the ethyl acetate phases combined; and the ethyl acetate phases washed with water, washed with brine, dried (sodium sulfate), and concentrated under reduced pressure to provide compound **1c** which was purified by silica gel chromatography using ethyl acetate/dichloromethane as the eluant. To compound **1c** (943 mg, 3.1 mmol) dissolved in 10 mL of ethyl acetate was added hydrochloric acid (4 M in dioxane, 7.7 mL, 30.7 mmol) and the resulting solution allowed to stir at room temperature for 16 hrs. The solution was then concentrated under reduced pressure, ethyl acetate added to the resulting residue, residual moisture removed by azeotropic distillation, and the remaining ethyl acetate removed under reduced pressure to provide the hydrochloride salt of **1d** which was stirred with diethyl ether and then filtered.

Compounds **ABA**, **AAV**, and **AAW** were prepared from the hydrochloride salt of **1d** using the following procedures:

Compound AAB: To the hydrochloride salt of **1d** (154 mg, 0.64 mmol) in anhydrous dichloromethane was added TEA (0.089 mL, 0.64 mmol) followed by 4-*tert*-butylphenyl isocyanate (112 mg, 0.64 mmol) and the resulting reaction mixture allowed to stir for 2 hours, after which the solvent was removed under reduced pressure to provide crude **ABA** which was purified by silica gel chromatography using ethyl acetate/dichloromethane as the eluant.

The identity of compounds **ABA** was confirmed by ¹H NMR. ¹H NMR (400 MHz, CDCl₃): δ 1.29 (s, 9H), 3.51 - 3.54 (m, 4H), 3.66 - 3.68 (m, 4H), 6.36 (bs, 1H), 7.16 (d, 2H), 7.27 (d, 2H).


Compound AAV: To the hydrochloride salt of 1d (154 mg, 0.64 mmol) in anhydrous DCM was added TEA (0.089 mL, 0.64 mmol) followed by 4-isopropyl phenyl isocyanate (103 mg, 0.64 mmol) and the resulting reaction mixture allowed to stir for 2 hours, after which the solvent was removed under reduced pressure to provide crude AAV which was purified by silica gel chromatography using ethyl acetate/dichloromethane as the eluant.

The identity of compounds AAV was confirmed by ¹H NMR. ¹H NMR (400 MHZ, CDCl₃): δ 1.23 (d, 6H), 2.84 - 2.90 (m, 1H), 3.53 - 3.55 (m, 4H), 3.65 - 3.67 (m, 4H), 6.31 (bs, 1H), 7.16 (d, 2H), 7.27 (d, 2H).

Compound AAW: To the hydrochloride salt of 1d (154 mg, 0.64 mmol) in anhydrous DCM was added TEA (0.089 mL, 0.64 mmol) followed by 4-trifluoromethyl phenyl isocyanate (120 mg, 0.64 mmol) and the resulting reaction mixture allowed to stir for 2 hours, after which the solvent was removed under reduced pressure to provide crude AAW which was purified by silica gel chromatography using ethyl acetate/dichloromethane as the eluant.

The identity of compounds AAW was confirmed by ¹H NMR. ¹H NMR (400 MHZ, CDCl₃): δ 3.55 - 3.57 (m, 4H), 3.71 - 3.73 (m, 4H), 6.41 (bs, 1H), 7.31 (d, 2H), 7.52 (d, 2H).

Thiadiazolylpiperazine Compounds of general formula:



wherein R is 4-methoxyphenyl, 4-trifluoromethoxyphenyl, 4-methylphenyl, or 4-chlorophenyl are made according to the procedure described above in Example 1 used to prepare compound ABA, except that the 4-*tert*-butylphenyl isocyanate is replaced with an equivalent of 4-methoxyphenyl isocyanate, 4-trifluoromethoxyphenyl isocyanate, 4-methylphenyl isocyanate, or 4-chlorophenyl isocyanate, respectively.

Example 2: Binding of the Thiadiazolylpiperazine Compounds to the Human VR1 Receptor


VR1 antagonism has been associated with the prevention of pain (See, e.g., R. Wrigglesworth and C. Walpole, "Capsaicin-like Agonists," *Drugs of the Future*, 23(5), pp. 531-538 (1998); S. Bevan P. McIntyre, "Vanilloid Receptors: Pivotal Molecules in Nociception," *Current Opinions in CPNS Investigational Drugs*, vol. 2, no. 2, pp. 178-185

(2000) and A. Szallash and P. Blumberg, "*Vanilloid (Capsaicin) Receptors and Mechanisms*," Pharmacology Reviews, vol. 51, no. 2, pp. 159-211 (1999)).

Methods for assaying compounds capable of inhibiting VR1 are well known to those skilled in the art, for example, those methods disclosed in U.S. Patent No. 6,239,267 to Duckworth *et al.*; U.S. Patent No. 6,406,908 to McIntyre *et al.*; or U.S. Patent No. 6,335,180 to Julius *et al.*

Typically, the Thiadiazolylpiperazine Compounds have an IC₅₀ value for inhibition of capsaicin-induced activation that is less than 25 μ M; in one embodiment, less than 1,000 nM; and in another embodiment, less than 500 nM, or an IC₅₀ value for inhibition of pH-induced inactivation that is less than 10 μ M; in one embodiment, less than 1,000 nM; and in another embodiment less than 500 nM.

The following Thiadiazolylpiperazine Compound inhibits capsaicin-induced activation and pH induced activation of VR1 as shown below:

Compound	Inhibition of capsaicin-induced activation IC ₅₀ (nM) Mean IC ₅₀ \pm SEC ^a (n) ^b	Inhibition of pH-induced activation IC ₅₀ (nM) Mean IC ₅₀ \pm SEC ^a (n) ^b
 <p style="text-align: center;">AAV</p>	374.4 \pm 233.4 (7)	668.3 \pm 265.1 (3)

^a SEM = standard deviation

^b n = number of experiments

5.3. Example 3: Binding of Thiadiazolepiperazine Compounds to mGluR5.

The following astrocyte assay can be used to demonstrate that

Thiadiazolylpiperazine Compounds bind to and modulate the activity of mGluR5.

Cell cultures: Primary glial cultures are prepared from cortices of Sprague-Dawley 18 days old embryos. The cortices are dissected and then dissociated by trituration. The resulting cell homogenate is plated onto poly-D-lysine precoated T175 flasks (BIOCOAT, commercially available from Becton Dickinson and Company Inc. of Franklin Lakes, NJ) in Dulbecco's Modified Eagle's Medium ("DMEM," pH 7.4), buffered with 25

mM HEPES, and supplemented with 15% fetal calf serum ("FCS," commercially available from Hyclone Laboratories Inc. of Omaha, NE), and incubated at 37° C and 5% CO₂. After 24 hours, FCS supplementation is reduced to 10%. On day six, oligodendrocytes and microglia are removed by strongly tapping the sides of the flasks. One day following this
5 purification step, secondary astrocyte cultures are established by subplating onto 96 poly-D-lysine precoated T175 flasks (BIOCOAT) at a density of 65,000 cells/well in DMEM and 10% FCS. After 24 hours, the astrocytes are washed with serum free medium and then cultured in DMEM, without glutamate, supplemented with 0.5% FCS, 20 mM HEPES, 10 ng/mL epidermal growth factor ("EGF"), 1 mM sodium pyruvate, and 1X
10 penicillin/streptomycin at pH 7.5 for 3 to 5 days at 37° C and 5% CO₂. The procedure allows the expression of the mGluR5 receptor by astrocytes, as demonstrated by S. Miller et al., *J. Neuroscience* 15(9):6103-6109 (1995).

Assay Protocol: After 3-5 days incubation with EGF, the astrocytes are washed with 127 mM NaCl, 5 mM KCl, 2 mM MgCl₂, 700 mM NaH₂PO₄, 2 mM CaCl₂, 5
15 mM NaHCO₃, 8 mM HEPES, 10 mM Glucose at pH 7.4 ("Assay Buffer") and loaded with the dye Fluo-4 (commercially available from Molecular Probes Inc. of Eugene, OR) using 0.1 mL of Assay Buffer containing Fluo-4 (3 mM final). After 90 minutes of dye loading, the cells are then washed twice with 0.2 mL Assay Buffer and resuspended in 0.1 mL of Assay Buffer. The plates containing the astrocytes are then transferred to a Fluorometric Imaging
20 Plate reader (commercially available from Molecular Devices Corporation of Sunnyvale, CA) for the assessment of calcium mobilization flux in the presence of glutamate and in the presence or absence of antagonist. After monitoring fluorescence for 15 seconds to establish a base line, DMSO solutions containing various concentrations of a Thiadiazolylpiperazine Compound diluted in Assay Buffer (0.05 mL of 4X dilutions for competition curves) are
25 added to the cell plate and fluorescence is monitored for 2 minutes. 0.05 mL of a 4X glutamate solution (agonist) is then added to each well to provide a final glutamate concentration in each well of 10 mM. Plate fluorescence is then monitored for an additional 60 seconds after agonist addition. The final DMSO concentration in the assay is 1.0%. In each experiment, fluorescence is monitored as a function of time and the data analyzed using
30 Microsoft Excel and GraphPad Prism. Dose-response curves are fit using a non-linear regression to determine IC₅₀ value. In each experiment, each data point is determined two times. The results demonstrate that Thiadiazolylpiperazine Compounds bind to and modulate the activity of mGluR5.

The following CHO assay can be used to demonstrate that
35 Thiadiazolylpiperazine Compounds bind to and modulate the activity of mGluR5.

40,000 CHO-rat mGluR5 cells/well are plated into 96 well plate (Costar 3409, Black, clear bottom, 96 well, tissue culture treated) for an overnight incubation in Dulbecco's Modified Eagle's Medium (DMEM, pH 7.4) supplemented with glutamine, 10% FBS, 1%

Pen/Strep, and 500ug/ml Geneticin. CHO-rat mGluR5 cells are washed and treated with Optimem medium and incubated for 1-4 hours prior to loading cells. Cell plates are then washed with loading buffer (127 mM NaCl, 5 mM KCl, 2 mM MgCl₂, 700 μ M Na H₂PO₄, 2 mM CaCl₂, 5 mM Na HCO₃, 8 mM Hepes, and 10 mM Glucose, pH 7.4) and then incubated with 3 μ M Fluo 4 (commercially available from Molecular probes Inc. of Eugene, OR) in 0.1 mL of loading buffer. After 90 minutes of dye loading, the cells are then washed twice with 0.2 mL loading buffer and resuspended in 0.1 mL loading buffer. The plates containing the CHO-rat mGluR5 cells are then transferred to a Fluorometric Imaging Plate Reader (FLIPR) (commercially available from Molecular Devices Corporation of Sunnyvale, CA) for the assessment of calcium mobilization flux in the presence of glutamate and in the presence or absence of test compounds. After monitoring fluorescence for 15 seconds to establish a baseline, DMSO solutions containing various concentrations of the test compound diluted in loading buffer (0.05 mL of 4X dilutions for the competition curves) are added to the cell plate and fluorescence is monitored for 2 minutes. 0.05 mL of 4X glutamate solution (agonist) is then added to each well to provide a final glutamate concentration in each well of 10 μ M. Plate fluorescence is then monitored for an additional 60 seconds after agonist addition. The final DMSO concentration in the assay is 1.0%. In each experiment, fluorescence is monitored as a function of time and the data analyzed using Microsoft Excel and GraphPad Prism. Dose-response curves are fit using a non-linear regression to determine IC₅₀ value. In each experiment each data point is determined two times.

5.4. Example 4: In Vivo Assays for Prevention or Treatment of Pain.

Test Animals: Each experiment uses rats weighing between 200-260 g at the start of the experiment. The rats are group-housed and have free access to food and water at all times, except prior to oral administration of a Thiadiazolylpiperazine Compound when food is removed for 16 hours before dosing. A control group acts as a comparison to rats treated with a Thiadiazolylpiperazine Compound. The control group is administered the carrier for the Thiadiazolylpiperazine Compound. The volume of carrier administered to the control group is the same as the volume of carrier and Thiadiazolylpiperazine Compound administered to the test group.

Acute Pain: To assess the actions of the Thiadiazolylpiperazine Compounds for the treatment or prevention of acute pain the rat tail flick test can be used. Rats are placed inside a cotton pouch and the tail exposed to a focused beam of radiant heat at a point 3 cm from the tip using a tail flick unit (Model 7360, commercially available from Ugo Basile of Italy). Tail flick latencies are defined as the interval between the onset of the thermal stimulus and the flick of the tail. Animals not responding within 15 seconds are removed from the tail flick unit and assigned a withdrawal latency of 15 seconds. Tail flick latencies are measured immediately before (pre-treatment) and 1, 3, and 6 hours following

administration of a Thiadiazolylpiperazine Compound. Data are expressed as tail flick latency(s) and the percentage of the maximal possible effect (% MPE), i.e., 15 seconds, is calculated as follows:

$$\%MPE = \frac{[(\text{post administration latency}) - (\text{pre - administration latency})]}{(15 \text{ s pre - administration latency})} \times 100\%$$

The rat tail flick test is described in F.E. D'Amour et al., "A Method for Determining Loss of Pain Sensation," *J. Pharmacol. Exp. Ther.* 72:74-79 (1941). The results demonstrate Thiadiazolylpiperazine Compounds that are useful for treating or preventing acute pain.

Acute pain can also be assessed by measuring the animal's response to noxious mechanical stimuli by determining the paw withdrawal threshold (PWT), as described below.

Inflammatory Pain: To assess the actions of the Thiadiazolylpiperazine Compounds for the treatment or prevention of inflammatory pain the Freund's complete adjuvant (FCA) model of inflammatory pain is used. FCA-induced inflammation of the rat hind paw is associated with the development of persistent inflammatory mechanical hyperalgesia and provides reliable prediction of the anti-hyperalgesic action of clinically useful analgesic drugs (L. Bartho et al., "Involvement of Capsaicin-sensitive Neurones in Hyperalgesia and Enhanced Opioid Antinociception in Inflammation," *Naunyn-Schmiedeberg's Archives of Pharmacology* 342:666-670 (1990)). The left hind paw of each animal is administered a 50 μ L intraplantar injection of 100% FCA. 24 hour post injection, the animal is assessed for response to noxious mechanical stimuli by determining the PWT, as described below. Rats are then administered a single injection of 1, 3, 10 or 30 mg/Kg of either a Thiadiazolylpiperazine Compound, 30 mg/Kg indomethacin or carrier. Responses to noxious mechanical stimuli are then determined 2, 4, 6, and 24 hours post administration. Percentage reversal of hyperalgesia for each animal is defined as:

$$\% \text{ Reversal} = \frac{[(\text{post administration PWT}) - (\text{pre - administration PWT})]}{(\text{Baseline pre - administration PWT})} \times 100\%$$

The results demonstrate Thiadiazolylpiperazine Compounds that are useful for treating or preventing inflammatory pain.

Neuropathic Pain: To assess the actions of the Thiadiazolylpiperazine Compounds for the treatment or prevention of neuropathic pain either the Seltzer model or the Chung model can be used.

In the Seltzer model, the partial sciatic nerve ligation model of neuropathic pain is used to produce neuropathic hyperalgesia in rats (Z. Seltzer et al., "A Novel

Behavioral Model of Neuropathic Pain Disorders Produced in Rats by Partial Sciatic Nerve Injury," *Pain* 43:205-218 (1990)). Partial ligation of the left sciatic nerve is performed under enflurane/O₂ inhalation anaesthesia. Following induction of anesthesia, the left thigh of the rat is shaved and the sciatic nerve exposed at high thigh level through a small incision and is carefully cleared of surrounding connective tissues at a site near the trochanter just distal to the point at which the posterior biceps semitendinosus nerve branches off of the common sciatic nerve. A 7-0 silk suture is inserted into the nerve with a 3/8 curved, reversed-cutting mini-needle and tightly ligated so that the dorsal 1/3 to 1/2 of the nerve thickness is held within the ligature. The wound is closed with a single muscle suture (7-0 silk) and a Michelle clip. Following surgery, the wound area is dusted with antibiotic powder. Sham-treated rats undergo an identical surgical procedure except that the sciatic nerve is not manipulated. Following surgery, animals are weighed and placed on a warm pad until they recover from anesthesia. Animals are then returned to their home cages until behavioral testing begins. The animal is assessed for response to noxious mechanical stimuli by determining PWT, as described below, immediately prior to and 1, 3, and 6 hours after drug administration for both the left rear paw and right rear paw of the animal. Percentage reversal of neuropathic hyperalgesia is defined as:

$$\% \text{ reversal} = 100 - [(\text{right pre-administration PWT} - \text{left post-administration PWT}) / (\text{right pre-administration PWT} - \text{left pre-administration PWT})] \times 100.$$

In the Chung model, the spinal nerve ligation model of neuropathic pain is used to produce mechanical hyperalgesia, thermal hyperalgesia and tactile allodynia in rats. Surgery is performed under isoflurane/O₂ inhalation anaesthesia. Following induction of anaesthesia a 3 cm incision is made and the left paraspinal muscles are separated from the spinous process at the L4 - S2 levels. The L6 transverse process is carefully removed with a pair of small rongeurs to identify visually the L4 - L6 spinal nerves. The left L5 (or L5 and L6) spinal nerve(s) is isolated and tightly ligated with silk thread. A complete hemostasis is confirmed and the wound is sutured using non-absorbable sutures, such as nylon sutures or stainless steel staples. Sham-treated rats undergo an identical surgical procedure except that the spinal nerve(s) is not manipulated. Following surgery animals are weighed, administered a subcutaneous (s.c.) injection of saline or ringers lactate, the wound area is dusted with antibiotic powder and they are kept on a warm pad until they recover from the anesthesia. Animals are then be returned to their home cages until behavioral testing begins. The animals are assessed for response to noxious mechanical stimuli by determining PWT, as described below, immediately prior to and 1, 3, and 5 hours after being administered a Thiadiazolylpiperazine Compound for both the left rear paw and right rear paw of the animal. The animal can also be assessed for response to noxious thermal stimuli or for tactile

allodynia, as described below. The Chung model for neuropathic pain is described in S.H. Kim, "An Experimental Model for Peripheral Neuropathy Produced by Segmental Spinal Nerve Ligation in the Rat," *Pain* 50(3):355-363 (1992). The results show demonstrate Thiadiazolylpiperazine Compounds that are useful for treating or preventing neuropathic pain.

Response to Mechanical Stimuli as an Assessment of Mechanical

Hyperalgesia: The paw pressure assay can be used to assess mechanical hyperalgesia. For this assay, hind paw withdrawal thresholds (PWT) to a noxious mechanical stimulus are determined using an analgesymeter (Model 7200, commercially available from Ugo Basile of Italy) as described in C. Stein, "Unilateral Inflammation of the Hindpaw in Rats as a Model of Prolonged Noxious Stimulation: *Alterations in Behavior and Nociceptive Thresholds*," *Pharmacology Biochemistry and Behavior* 31:451-455 (1988). The maximum weight that can be applied to the hind paw is set at 250 g and the end point is taken as complete withdrawal of the paw. PWT is determined once for each rat at each time point and only the affected (ipsilateral) paw is tested.

Response to Thermal Stimuli as an Assessment of Thermal Hyperalgesia:

The plantar test can be used to assess thermal hyperalgesia. For this test, hind paw withdrawal latencies to a noxious thermal stimulus are determined using a plantar test apparatus (commercially available from Ugo Basile of Italy) following the technique described by K. Hargreaves et al., "A New and Sensitive Method for Measuring Thermal Nociception in Cutaneous Hyperalgesia," *Pain* 32(1):77-88 (1988). The maximum exposure time is set at 32 seconds to avoid tissue damage and any directed paw withdrawal from the heat source is taken as the end point. Three latencies are determined at each time point and averaged. Only the affected (ipsilateral) paw is tested.

Assessment of Tactile Allodynia:

To assess tactile allodynia, rats are placed in clear, plexiglass compartments with a wire mesh floor and allowed to habituate for a period of at least 15 minutes. After habituation, a series of von Frey monofilaments are presented to the plantar surface of the left (operated) foot of each rat. The series of von Frey monofilaments consists of six monofilaments of increasing diameter, with the smallest diameter fiber presented first. Five trials are conducted with each filament with each trial separated by approximately 2 minutes. Each presentation lasts for a period of 4-8 seconds or until a nociceptive withdrawal behavior is observed. Flinching, paw withdrawal or licking of the paw are considered nociceptive behavioral responses.

5.5. Example 5: In Vivo Assays for Prevention or Treatment of Anxiety.

The elevated plus maze test or the shock-probe burying test can be used to assess the anxiolytic activity of Thiadiazolylpiperazine Compounds in rats or mice.

The Elevated Plus Maze Test: The elevated plus maze consists of a platform with 4 arms, two open and two closed (50x10x50 cm enclosed with an open roof). Rats (or mice) are placed in the center of the platform, at the crossroad of the 4 arms, facing one of the closed arms. Time spent in the open arms vs the closed arms and number of open arm entries during the testing period are recorded. This test is conducted prior to drug administration and again after drug administration. Test results are expressed as the mean time spent in open arms and the mean number of entries into open arms. Known anxiolytic drugs increase both the time spent in open arms and number of open arm entries. The elevated plus maze test is described in D. Treit, "Animal Models for the Study of Anti-anxiety Agents: A Review," *Neuroscience & Biobehavioral Reviews* 9(2):203-222 (1985).

The Shock-Probe Burying Test: For the shock-probe burying test the testing apparatus consists of a plexiglass box measuring 40x30x40 cm, evenly covered with approximately 5 cm of bedding material (odor absorbent kitty litter) with a small hole in one end through which a shock probe (6.5 cm long and 0.5 cm in diameter) is inserted. The plexiglass shock probe is helically wrapped with two copper wires through which an electric current is administered. The current is set at 2 mA. Rats are habituated to the testing apparatus for 30 min on 4 consecutive days without the shock probe in the box. On test day, rats are placed in one corner of the test chamber following drug administration. The probe is not electrified until the rat touches it with its snout or fore paws, at which point the rat receives a brief 2 mA shock. The 15 min testing period begins once the rat receives its first shock and the probe remains electrified for the remainder of the testing period. The shock elicits burying behavior by the rat. Following the first shock, the duration of time the rat spends spraying bedding material toward or over the probe with its snout or fore paws (burying behavior) is measured as well as the number of contact-induced shocks the rat receives from the probe. Known anxiolytic drugs reduce the amount of burying behavior. In addition, an index of the rat's reactivity to each shock is scored on a 4 point scale. The total time spent immobile during the 15 min testing period is used as an index of general activity. The shock-probe burying test is described in D. Treit, 1985, *supra*. The results demonstrate Thiadiazolylpiperazine Compounds that are useful for treating or preventing anxiety.

5.6. Example 6: In Vivo Assays for Prevention or Treatment of an Addictive Disorder.

The condition place preference test or drug self-administration test can be used to assess the ability of Thiadiazolylpiperazine Compounds to attenuate the rewarding properties of known drugs of abuse.

The Condition Place Preference Test: The apparatus for the conditioned place preference test consists of two large compartments (45x45x30 cm) made of wood with a plexiglass front wall. These two large compartments are distinctly different. Doors at the back of each large compartment lead to a smaller box (36x18x20 cm) box made of wood,

painted grey, with a ceiling of wire mesh. The two large compartments differ in terms of shading (white vs black), level of illumination (the plexiglass door of the white compartment is covered with aluminum foil except for a window of 7x7 cm), texture (the white compartment has a 3 cm thick floor board (40x40 cm) with nine equally spaced 5 cm diameter holes and the black has a wire mesh floor), and olfactory cues (saline in the white compartment and 1 mL of 10% acetic acid in the black compartment). On habituation and testing days, the doors to the small box remain open, giving the rat free access to both large compartments.

The first session that a rat is placed in the apparatus is a habituation session and entrances to the smaller grey compartment remain open giving the rat free access to both large compartments. During habituation, rats generally show no preference for either compartment. Following habituation, rats are given 6 conditioning sessions. Rats are divided into 4 groups: carrier pre-treatment + carrier (control group), 2 Thiadiazolylpiperazine Compound pre-treatment + carrier, carrier pre-treatment + morphine, 2 Thiadiazolylpiperazine Compound pre-treatment + morphine. During each conditioning session the rat is injected with one of the drug combinations and confined to one compartment for 30 min. On the following day, the rat receives a carrier + carrier treatment and is confined to the other large compartment. Each rat receives three conditioning sessions consisting of 3 drug combination-compartment and 3 carrier-compartment pairings. The order of injections and the drug/compartment pairings are counterbalanced within groups. On the test day, rats are injected prior to testing (30 min to 1 hour) with either morphine or carrier and the rat is placed in the apparatus, the doors to the grey compartment remain open and the rat is allowed to explore the entire apparatus for 20 min. The time spent in each compartment is recorded. Known drugs of abuse increase the time spent in the drug-paired compartment during the testing session. If the Thiadiazolylpiperazine Compound blocks the acquisition of morphine conditioned place preference (reward), there will be no difference in time spent in each side in rats pre-treated with a Thiadiazolylpiperazine Compound and the group will not be different from the group of rats that was given carrier + carrier in both compartments. Data will be analyzed as time spent in each compartment (drug combination-paired vs carrier-paired). Generally, the experiment is repeated with a minimum of 3 doses of a Thiadiazolylpiperazine Compound.

The Drug Self-Administration Test: The apparatus for the drug self-administration test is a standard commercially available operant conditioning chamber. Before drug trials begin rats are trained to press a lever for a food reward. After stable lever pressing behavior is acquired, rats are tested for acquisition of lever pressing for drug reward. Rats are implanted with chronically indwelling jugular catheters for i.v. administration of compounds and are allowed to recover for 7 days before training begins. Experimental sessions are conducted daily for 5 days in 3 hour sessions. Rats are trained to self-administer

a known drug of abuse, such as morphine. Rats are then presented with two levers, an "active" lever and an "inactive" lever. Pressing of the active lever results in drug infusion on a fixed ratio 1 (FR1) schedule (*i.e.*, one lever press gives an infusion) followed by a 20 second time out period (signaled by illumination of a light above the levers). Pressing of the inactive lever results in infusion of excipient. Training continues until the total number of morphine infusions stabilizes to within $\pm 10\%$ per session. Trained rats are then used to evaluate the effect of Thiadiazolylpiperazine Compounds pre-treatment on drug self-administration. On test day, rats are pre-treated with a Thiadiazolylpiperazine Compound or excipient and then are allowed to self-administer drug as usual. If the Thiadiazolylpiperazine Compound blocks the rewarding effects of morphine, rats pre-treated with the Thiadiazolylpiperazine Compound will show a lower rate of responding compared to their previous rate of responding and compared to excipient pre-treated rats. Data is analyzed as the change in number of drug infusions per testing session (number of infusions during test session – number of infusions during training session). The results show that Thiadiazolylpiperazine Compounds are useful for treating or preventing an addictive disorder.

5.7. Example 7: Functional Assay for Characterizing mGluR 1 Antagonistic Properties.

Functional assays for the characterization of mGluR1 antagonistic properties are well known in the art. For example, the following procedure can be used.

cDNA encoding rat mGluR1a receptor is obtained from, *e.g.*, Prof. S. Nakanishi (Kyoto, Japan). It is transiently transfected into HEK-EBNA cells using a procedure described by Schlaeger et al., *New Dev. New Appl. Anim. Cell Techn., Proc. ESACT Meet.*, (1998), 105-112 and 117-120. $[Ca^{2+}]$ measurements are performed on mGluR1a transfected HEK-EBNA cells after incubation of the cells with Fluo-3 AM (0.5 μ M final concentration) for 1 hour at 37°C followed by 4 washes with assay buffer (DMEM supplemented with Hank's salt and 20 mM HEPES). $[Ca^{2+}]$ measurements are done using a fluorometric imaging plate reader, *e.g.*, FLIPR from Molecular Devices Corporation, La Jolla, CA. 10 μ M glutamate as agonist is used to evaluate the potency of the antagonists.

Increasing concentrations of antagonists are applied to the cells 5 minutes prior to application of the agonist. The inhibition (antagonists) curves are fitted with appropriate software, for example, the four-parameter logistic equation giving IC_{50} and Hill coefficient using the iterative nonlinear curve fitting software Origin from Microcal Software Inc., Northampton, MA.


The present invention is not to be limited in scope by the specific embodiments disclosed in the examples which are intended as illustrations of a few aspects of the invention and any embodiments that are functionally equivalent are within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and

described herein will become apparent to those skilled in the art and are intended to fall within the scope of the appended claims.

A number of references have been cited, the entire disclosures of which are incorporated herein by reference.

What is claimed is:

1. A compound of formula:



(I)

or a pharmaceutically acceptable salt thereof, wherein:

R^1 is methyl or halo;

R^3 is:

- 10 (a) $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, $-(C_2-C_6)alkynyl$, or $-(CH_2)_nOR^6$,
which is unsubstituted or substituted with one or more of $-CN$, $-OH$, $-halo$, $-NO_2$, $-CH=NR^6$, or $-NR^6OH$; or
- (b) $-H$, $-halo$, $-NO_2$, $-CN$, or $-NH_2$;

R^4 is $-H$;

- 15 R^5 is:

- (a) $-(C_1-C_6)alkyl$, $-(C_2-C_6)alkenyl$, or $-(C_2-C_6)alkynyl$; or
- (b) $-(C_3-C_8)cycloalkyl$, $-(C_5-C_8)cycloalkenyl$, $-phenyl$, $-naphthyl$, $-(C_{14})aryl$, $-(C_3-C_7)heterocycle$, $-(C_1-C_6)alkylphenyl$, $-(C_1-C_6)alkylnaphthyl$, $-(C_1-C_6)alkyl(C_{14})aryl$, $-(C_1-C_6)alkyl(C_3-C_7)heterocycle$,
20 $-(C_2-C_6)alkenylphenyl$, $-(C_2-C_6)alkenylphenyl$, $-(C_2-C_6)alkenyl(C_{14})aryl$,
 $-(C_2-C_6)alkenyl(C_3-C_7)heterocycle$, $-(C_2-C_6)alkynyl(C_3-C_8)cycloalkyl$,
 $-(C_2-C_6)alkynyl(C_5-C_8)cycloalkenyl$, $-(C_2-C_6)alkynylphenyl$,
 $-(C_2-C_6)alkylnaphthyl$, $-(C_2-C_6)alkynyl(C_{14})aryl$, or

-(C₂-C₆)alkynyl(C₃-C₇)heterocycle, which is unsubstituted or substituted with one or more R⁷ groups;

each R⁶ is independently -H, -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -(C₃-C₈)cycloalkyl, -(C₅-C₈)cycloalkenyl, -phenyl, -(C₃-C₅)heterocycle, C(halo)₃, or CH(halo)₂;

each R⁷ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -(CH₂)_n(C₃-C₈)cycloalkyl, -(CH₂)_n(C₅-C₈)cycloalkenyl, -(CH₂)_nphenyl, -(CH₂)_nSR⁶, -(CH₂)_nCH(halo)₂, -(CH₂)_nC(halo)₃, -halo, -(CH₂)_nOR⁶; and

n is an integer ranging from 0 to 2.

2. The compound of claim 1, wherein R³ and R⁴ is H; and R⁵ is phenyl which is unsubstituted or substituted with a -(C₁ - C₆)alkyl.

3. The compound of claim 2, wherein R¹ is a halogen.

4. The compound of claim 3, wherein R¹ is chloro.


5. The compound of claim 2, wherein R¹ is a methyl.

6. The compound of claim 2, wherein the phenyl group is substituted in the 4-position.

7. The compound of claim 6, wherein the -(C₁-C₆)alkyl group is a *t*-butyl group.

8. The compound of claim 1, wherein R¹ is methyl.

9. A compound of formula:



(II)

or a pharmaceutically acceptable salt thereof, wherein:

R¹ is methyl or halo;

R³ is:

15 (a) -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, or -(CH₂)_nOR⁶,
which R³ and R⁴ are each independently unsubstituted or substituted with one
or more of -CN, -OH, -halo, -NO₂, -CH=NR⁶, or -NR⁶OH; or

(b) -H, -halo, -NO₂, -CN, or -NH₂;

R⁴ is: -H;

R⁵ is:

20 (a) -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, or -(C₂-C₆)alkynyl; or

(b) -(C₃-C₈)cycloalkyl, -(C₅-C₈)cycloalkenyl, -phenyl, -naphthyl, -
(C₁₄)aryl, -(C₃-C₇)heterocycle, -(C₁-C₆)alkylphenyl, -(C₁-C₆)alkylnaphthyl, -
(C₁-C₆)alkyl(C₁₄)aryl, -(C₁-C₆)alkyl(C₃-C₇)heterocycle,
25 -(C₂-C₆)alkenylphenyl, -(C₂-C₆)alkenylnaphthyl, -(C₂-C₆)alkenyl(C₁₄)aryl,
-(C₂-C₆)alkenyl(C₃-C₇)heterocycle, -(C₂-C₆)alkynyl(C₃-C₈)cycloalkyl,
-(C₂-C₆)alkynyl(C₅-C₈)cycloalkenyl, -(C₂-C₆)alkynylphenyl,
-(C₂-C₆)alkynylnaphthyl, -(C₂-C₆)alkynyl(C₁₄)aryl, or
-(C₂-C₆)alkynyl(C₃-C₇)heterocycle, which is unsubstituted or substituted with
one or more R⁷ groups;

30 each R⁶ is independently -H, -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl,
-(C₃-C₈)cycloalkyl, -(C₅-C₈)cycloalkenyl, -phenyl, -(C₃-C₅)heterocycle, C(halo)₃, or
CH(halo)₂;

each R⁷ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl,
-(CH₂)_n(C₃-C₈)cycloalkyl, -(CH₂)_n(C₅-C₈)cycloalkenyl, -(CH₂)_nphenyl, -(CH₂)_nSR⁶,
35 -(CH₂)_nCH(halo)₂, -(CH₂)_nC(halo)₃, -halo, -(CH₂)_nOR⁶; and

n is an integer ranging from 0 to 2.

10. The compound of claim 9, wherein R³ and R⁴ is H, and R⁵ is phenyl which is
unsubstituted or substituted with a -(C₁ - C₆)alkyl.

11. The compound of claim 10, wherein R¹ is a halogen.

12. The compound of claim 11, wherein R¹ is chloro.

13. The compound of claim 10, wherein R¹ is a methyl.
14. The compound of claim 10, wherein the phenyl group is substituted in the 4-position.
15. The compound of claim 14, wherein the -(C₁-C₆)alkyl group is a *t*-butyl group.
16. The compound of claim 9, wherein R¹ is methyl.
17. A pharmaceutical composition comprising the compound or a pharmaceutically acceptable salt of the compound of claim 1 and a pharmaceutically acceptable carrier or excipient.
18. The composition of claim 17, further comprising an opioid analgesic.
19. The composition of claim 17, further comprising a non-opioid analgesic.
20. The composition of claim 17, further comprising an anti-emetic agent.
21. A pharmaceutical composition comprising a compound or a pharmaceutically acceptable salt of the compound of claim 9 and a pharmaceutically acceptable carrier or excipient.
22. The composition of claim 21, further comprising an opioid analgesic.
23. The composition of claim 21, further comprising a non-opioid analgesic.
24. The composition of claim 21, further comprising an anti-emetic agent.
25. A method of treating pain in a patient, comprising administering to a patient in need thereof an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 1.
26. The method of claim 25, further comprising administering an effective amount of an opioid analgesic.
27. The method of claim 25, further comprising administering an effective amount of a non-opioid analgesic.

28. The method of claim 25, further comprising administering an effective amount of an anti-emetic agent.

29. A method of treating pain in a patient, comprising administering to a patient in need thereof an effective amount of a compound or a pharmaceutically acceptable salt of the compound of claim 9.

30. The method of claim 29, further comprising administering an effective amount of an opioid analgesic.

31. The method of claim 29, further comprising administering an effective amount of a non-opioid analgesic.

32. The method of claim 29, further comprising administering an effective amount of an anti-emetic agent.

33. A method for inhibiting VR1 receptor function in a cell comprising contacting a cell capable of expressing VR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 1.

34. A method for inhibiting VR1 receptor function in a cell comprising contacting a cell capable of expressing with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 9.

35. A method for treating depression in an animal, comprising administering to an animal in need thereof an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 1.

36. The method of claim 35, further comprising administering an effective amount of an antidepressant.

37. A method for treating depression in an animal, comprising administering to an animal in need thereof an effective amount of a compound or a pharmaceutically acceptable salt of the compound of claim 9.

38. The method of claim 37, further comprising administering an effective amount of an antidepressant.

39. A method for inhibiting mGluR1 receptor function in a cell comprising contacting a cell capable of expressing mGluR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 1.

40. A method for inhibiting mGluR1 receptor function in a cell comprising contacting a cell capable of expressing mGluR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 9.

41. A method for treating anxiety in an animal, comprising administering to an animal in need thereof an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 1.

42. The method of claim 41, further comprising administering an effective amount of an anti-anxiety agent.


43. A method for treating anxiety in an animal, comprising administering to an animal in need thereof an effective amount of a compound or a pharmaceutically acceptable salt of the compound of claim 9.

44. The method of claim 37, further comprising administering an effective amount of an anti-anxiety agent.

45. A method for inhibiting mGluR5 receptor function in a cell comprising contacting a cell capable of expressing mGluR5 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 1.

46. A method for inhibiting mGluR5 receptor function in a cell comprising contacting a cell capable of expressing mGluR5 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 9.

47. A compound of formula :



or a pharmaceutically acceptable salt thereof, wherein:

X is O or S;

R¹ is -CH₃ or -halo;

R³ is:

(a) -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, or -(CH₂)ₙOR⁶, which is unsubstituted or substituted with one or more of -CN, -OH, -halo, -NO₂, -CH=NR⁶, or -NR⁶OH; or

(b) -H, -halo, -NO₂, -CN, or -NH₂;

R⁴ is -H;

each R⁸ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -(CH₂)ₙ(C₃-C₈)cycloalkyl, -(CH₂)ₙ(C₅-C₈)cycloalkenyl, -(CH₂)ₙphenyl, -(CH₂)ₙSR⁶, -(CH₂)ₙCH(halo)₂, -(CH₂)ₙC(halo)₃, -halo, -(CH₂)ₙOR⁶;

each R⁶ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -phenyl, -naphthyl, or -(C₁₄)aryl;


Ar is -phenyl, -naphthyl, -(C₁₄)aryl, -(C₃-C₈)cycloalkyl or -(C₅-C₈)cycloalkenyl, which is unsubstituted or substituted with one or more R⁷ groups;

each R⁷ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -(CH₂)ₙ(C₃-C₈)cycloalkyl, -(CH₂)ₙ(C₅-C₈)cycloalkenyl, -(CH₂)ₙphenyl, -(CH₂)ₙSR⁶, -(CH₂)ₙCH(halo)₂, -(CH₂)ₙC(halo)₃, -halo, -(CH₂)ₙOR⁶;

n is an integer ranging from 0 to 2; and

p is an integer ranging from 0 to 2.

48. The compound of claim 47, wherein X is O.
49. The compound of claim 48, wherein R¹ is methyl.
50. The compound of claim 49, wherein Ar is phenyl.
51. The compound of claim 50, wherein the phenyl is substituted in the 4-position with a -(C₁-C₆)alkyl group.
52. The compound of claim 51, wherein the -(C₁-C₆)alkyl group is a *t*-butyl group.
53. The compound of claim 48, wherein R¹ is -halo.
54. The compound of claim 53, wherein Ar is phenyl.
55. The compound of claim 54, wherein the phenyl is substituted in the 4-position with a -(C₁-C₆)alkyl group.
56. The compound of claim 55, wherein the -(C₁-C₆)alkyl group is a *t*-butyl group.
57. The compound of claim 56, wherein R¹ is chloro.
58. The compound of claim 47, wherein X is S.
59. A compound of formula:



(IV)

25 or a pharmaceutically acceptable salt thereof, wherein:

X is O or S;

R¹ is -CH₃ or -halo;

R³ is:

30 (a) -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, or -(CH₂)_nOR⁶,
which is unsubstituted or substituted with one or more of -CN, -OH, -halo, -
NO₂, -CH=NR⁶, or -NR⁶OH; or

(b) -H, -halo, -NO₂, -CN, or -NH₂;

R⁴ is -H;

35 each R⁸ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl,
-(CH₂)_n(C₃-C₈)cycloalkyl, -(CH₂)_n(C₅-C₈)cycloalkenyl, -(CH₂)_nphenyl, -(CH₂)_nSR⁶,
-(CH₂)_nCH(halo)₂, -(CH₂)_nC(halo)₃, -halo, -(CH₂)_nOR⁶;

each R⁶ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl, -
phenyl, -naphthyl, or -(C₁₄)aryl;

40 Ar is -phenyl, -naphthyl, -(C₁₄)aryl, -(C₃-C₈)cycloalkyl or -(C₅-C₈)cycloalkenyl,
which is unsubstituted or substituted with one or more R⁷ groups;

each R⁷ is independently -(C₁-C₆)alkyl, -(C₂-C₆)alkenyl, -(C₂-C₆)alkynyl,
-(CH₂)_n(C₃-C₈)cycloalkyl, -(CH₂)_n(C₅-C₈)cycloalkenyl, -(CH₂)_nphenyl, -(CH₂)_nSR⁶,
-(CH₂)_nCH(halo)₂, -(CH₂)_nC(halo)₃, -halo, -(CH₂)_nOR⁶;

n is an integer ranging from 0 to 2; and

45 p is an integer ranging from 0 to 2.

60. The compound of claim 59, wherein X is O.

61. The compound of claim 60, wherein R¹ is methyl.

62. The compound of claim 61, wherein Ar is phenyl.

63. The compound of claim 62, wherein the phenyl is substituted in the 4-position with a -(C₁-C₆)alkyl group.

64. The compound of claim 63, wherein the -(C₁-C₆)alkyl group is a *t*-butyl group.

65. The compound of claim 60, wherein R¹ is -halo.

66. The compound of claim 65, wherein Ar is phenyl.

67. The compound of claim 66, wherein the phenyl is substituted in the 4-position with a -(C₁-C₆)alkyl group.

68. The compound of claim 66, wherein the -(C₁-C₆)alkyl group is a *t*-butyl group.

69. The compound of claim 68, wherein R¹ is chloro.

70. The compound of claim 59, wherein X is S.

71. A pharmaceutical composition comprising an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47 and a pharmaceutically acceptable carrier or excipient.

72. The composition of claim 71, further comprising an opioid analgesic.

73. The composition of claim 71, further comprising a non-opioid analgesic.

74. The composition of claim 71, further comprising an anti-emetic agent.

75. A pharmaceutical composition comprising an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 59 and a pharmaceutically acceptable carrier or excipient.

76. The composition of claim 75, further comprising an opioid analgesic.

77. The composition of claim 75, further comprising a non-opioid analgesic.
78. The composition of claim 75, further comprising an anti-emetic agent.
79. A method for treating pain in an animal, comprising administering to an animal in need thereof an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47.
80. The method of claim 79, further comprising administering an effective amount of an opioid analgesic.
81. The method of claim 79, further comprising administering an effective amount of a non-opioid analgesic.
82. The method of claim 79, further comprising administering an effective amount of an anti-emetic agent.
83. A method for treating pain in an animal, comprising administering to an animal in need thereof an effective amount of a compound or a pharmaceutically acceptable salt of the compound of claim 59.
84. The method of claim 83, further comprising administering an effective amount of an opioid analgesic.
85. The method of claim 83, further comprising administering an effective amount of a non-opioid analgesic.
86. The method of claim 83, further comprising administering an effective amount of an anti-emetic agent.
87. A method for inhibiting VR1 receptor function in a cell comprising contacting a cell capable of expressing VR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47.
88. A method for inhibiting VR1 receptor function in a cell comprising contacting a cell capable of expressing VR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 59.

89. A method for treating depression in an animal, comprising administering to an animal in need thereof an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47.

90. The method of claim 89, further comprising administering an effective amount of an antidepressant.

91. A method for treating depression in an animal, comprising administering to an animal in need thereof an effective amount of a compound or a pharmaceutically acceptable salt of the compound of claim 59.

92. The method of claim 91, further comprising administering an effective amount of an antidepressant.

93. A method for inhibiting mGluR1 receptor function in a cell comprising contacting a cell capable of expressing mGluR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47.

94. A method for inhibiting mGluR1 receptor function in a cell comprising contacting a cell capable of expressing mGluR1 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 59.

95. A method for treating anxiety in an animal, comprising administering to an animal in need thereof an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47.

96. The method of claim 95, further comprising administering an effective amount of an anti-anxiety agent.

97. A method for treating anxiety in an animal, comprising administering to an animal in need thereof an effective amount of a compound or a pharmaceutically acceptable salt of the compound of claim 59.

98. The method of claim 97, further comprising administering an effective amount of an anti-anxiety agent.

99. A method for inhibiting mGluR5 receptor function in a cell comprising contacting a cell capable of expressing mGluR5 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 47.

100. A method for inhibiting mGluR5 receptor function in a cell comprising contacting a cell capable of expressing mGluR5 with an effective amount of the compound or a pharmaceutically acceptable salt of the compound of claim 59.

INTERNATIONAL SEARCH REPORT

Internatly Application No

PCT/US 03/06107

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/04 A61K31/496 A61P25/22 A61P25/24 A61P29/00
 //(C07D417/04,241:00,285:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 51999 A (AMERICAN HOME PROD) 8 September 2000 (2000-09-08) Scheme 1 (interm. I, III and IV), schemes 2 and 3 (interm. VIII and IX). The intermediate tert-butyl ester of example 1, interm. I and II of example 4, IV and V of example 7 and X of example 11. abstract; claims 13,14 ----	1-100
A	EP 0 438 230 A (MERCK SHARP & DOHME) 24 July 1991 (1991-07-24) example 96 page 9, line 14 - line 15; claim 6 -----	1-100
A	WO 00 35915 A (NAKAI KAZUO ; TAKE KAZUHIKO (JP); AZAMI HIDENORI (JP); EIKYU YOSHIT) 22 June 2000 (2000-06-22) abstract; claim 1 -----	1-100

☐ Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

18 June 2003

Date of mailing of the international search report

27/06/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Papathoma, S

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 03/06107

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 25-46 and 79-100 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

In relation on patent family members

Internati Application No

PCT/US 03/06107

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0051999	A	08-09-2000	AU 3247800 A	21-09-2000
			CA 2361179 A1	08-09-2000
			CN 1349520 T	15-05-2002
			EP 1161424 A1	12-12-2001
			JP 2002538152 A	12-11-2002
			WO 0051999 A1	08-09-2000
EP 0438230	A	24-07-1991	AT 152110 T	15-05-1997
			AU 6944091 A	25-07-1991
			CA 2034189 A1	18-07-1991
			CN 1053429 A	31-07-1991
			CS 9100086 A2	15-09-1991
			DE 69125731 D1	28-05-1997
			DE 69125731 T2	27-11-1997
			EP 0438230 A2	24-07-1991
			FI 910228 A	18-07-1991
			IE 910139 A1	17-07-1991
			JP 6100558 A	12-04-1994
			NO 910187 A	18-07-1991
			PT 96477 A	15-10-1991
WO 0035915	A	22-06-2000	AU 1683700 A	03-07-2000
			BR 9917047 A	30-07-2002
			CA 2354875 A1	22-06-2000
			CN 1334812 T	06-02-2002
			CZ 20012130 A3	17-10-2001
			EP 1140924 A1	10-10-2001
			HU 0203651 A2	28-02-2003
			WO 0035915 A1	22-06-2000
			JP 2002532499 T	02-10-2002
			TR 200101649 T2	22-10-2001

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKewed/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINEs OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)